




2016

ABSTRACT POSTER "IV INTERNATIONAL COURSE OF ORBITOPLASTIC SURGERY IN AMALFI COAST"

NEUROFTALMOLOGIA

OFTALMOPLASTIA



ESTÈTIQUE

SURGERY

**ABSTRACT POSTER  
"IV INTERNATIONAL COURSE  
OF ORBITOPLASTIC SURGERY  
IN AMALFI COAST"**

*Edited by*  
**Luigi Colangelo**

ISBN 979-12-200-1977-4



9 791220 019774

OCTOBER 7 - 8 th 2016  
PRAIANO - SALERNO - ITALY

**IV INTERNATIONAL COURSE  
OF ORBITOPLASTIC SURGERY**

**OCTOBER 7-8 th 2016 PRAIANO  
SALERNO - ITALY**

**ORGANIZER:  
Luigi COLANGELO**

**SCIENTIFIC COMMITTEE:**

**President: Daniel Weil**

Luigi Colangelo

Ana Rosa Pimentel

Nicola Rosa

Elio Trotta

**COMMITTEE FOR POSTERS**

**President: LUCA CIMINO**

Giovanni Cennamo

Lino Guarracino

Eduardo Rubin

Jose Luis Tovilla



## TABLE OF CONTENTS

Preface.....	1
‘Francesco Colangelo Award Lecture’ .....	2
<b>N°1 Evaluation of the radial peripapillary capillary network in congenital optic disc anomalies with optical coherence tomography angiography</b>	
Gilda Cennamo MD, Claudia Rossi MD, Pasquale RuggeroMD, and Giovanni Cennamo MD.....	5
<b>N°2 Leber hereditary optic neuropathy: a case report</b>	
Gianluca Baldi, Giovanni Leo, Maurizio Cioffi, Luca Barile, Assunta Rosaria Cascone, Alfonso D’Alessandro.....	6
<b>N°3 First italian family with autosomal recessive retinal dystrophy caused by MERTK mutation (P38)</b>	
Gallenga CE, Lanciotti S, Lattanzi D, Bolletta E, Parmeggiani F, Perri P, Giovannini A, Novelli G, Brancati F. ....	7
<b>N°4 Improving VA in Stargardt disease using Gabor patch technology: a retrospective study</b>	
Mincarelli C, Gallenga CE, Colangelo L, Lobefalo L, Rapinese M, Di Carlo PL, Nuri M, Perri P, Parmeggiani F.....	10
<b>N°5 Exenteration of orbit in a tertiary eye hospital - indication and outcomes</b>	
Silvana A. Schellini, Alicia Galindo-Ferreiro, Rajiv Khandekar, Patricia M. Akaishi, Antonio Augusto V. Cruz, Yerena Muiños .....	12
<b>N°6 Expansive hydrogel pellets in the treatment of post enucleation/evisceration socket syndrome</b>	
Delmas Juliette, Lossouran Adrien, Adenis Jean-Paul, Robert Pierre-Yves.....	15
<b>N°7 Our experience with an automaded photo-ophthalmic device for measuring MRD</b>	
Simon Barnard & Giuseppe Reppucci .....	16
<b>N°8 Surgical management of residual brow suspension ptosis and severe lagophthalmos</b>	
Dr.Michel Tazartes, Dr.Sorinela Roata, Dr. Saad Mouine.....	17
<b>N° 9 Orbital and adnexa involvement in angiolymphod hyperplasia with epheliphilia</b>	
Alicia Galindo-Ferreiro, Hind M Alkatan, Sahar Elkhamary, Azza Maktabi, Silvana Schellini, Antonio Augusto Cruz. Patrica Akaishi.....	18

<b>N° 10 Orbital tumor in childhood, un unsuspected presentation</b>	
Avila Jorge Ezequiel.....	21
<b>N°11 Spontaneous preoperative rupture of dermoid cysts: clinicopathological and radiological features</b>	
Alicia Galindo-Ferreiro, Patricia Akaishi, Yerena Muiños-Diaz, Silvana Schellini, Augusto Cruz, Sahar Elkhamary, Hind Alkhatan, Rajiv Khandekar .....	23
<b>N°12 Endoscopic primary dacryocystorhinostomy: are silicone tubes needed?</b>	
M. Cavaliere, M.D.; G. Concilio, M.D.; F.M. Folino, M.D.; M. Iemma, M.D.....	26
<b>N°13 Results of probing under endoscopic guidance in children foury-eight months and older</b>	
Silvana Schellini MD, Alicia Galindo-Ferreiro, Rajiv Khandekar, Patricia Mitiko Akaishi, Augusto Cruz .....	28
<b>N°14 Temporary medial upper eyelid lagophthalmos after external dacryocystorhinostomy</b>	
Ivan O. Haefliger, M.D., FEBO, Otmar Meienberg, M.D. and Ana Rosa Pimentel de Figueiredo, M.D., Ph.D .....	31
<b>N°15 Differents forms of conjunctival and eyelid lymphomas</b>	
Cam.acho Belén, Ignacio Racana, Tula Romina, Billagra Alejandra, Weil Daniel.....	32
<b>N°16 Sebaceous carcinoma of the eyelid with pagetoid infiltration of the conjunctiva: two case reports</b>	
Ricardo Díaz Céspedes, Gloria Gornals Montés, Diamar Pardo Lopez Enrique España Gregori, David Salom Alonso .....	34
<b>N°17 Total reconstruction Functional of the upper and lower eyelid using combined flaps and free grafts</b>	
Torres Ana Marina. MD Coauthor: Dolmetsch Angela M, MD .....	36
<b>N°18 Transconjunctival fat reposition without pedicle in lower blepharoplasty</b>	
Torres, A. Marina .....	37
<b>N°19 Orbital mucocele</b>	
Arce.Jessica, Avila.Jorge, Billagra.Alejandra, Camacho. Belén, Gasparini.Cecilia, Magnano. Paula, Racana. Ignacio, Tula. Romina, Weil. Daniel.....	38
<b>N°20 Orbital hemangiopericytoma</b>	
Cynthia Anagua, Cecilia Gasparini, Sabrina Bagdadi, Martin Rojo, Daniel Weil .....	39
<b>N° 21 The Garden of Minerva</b>	

Billagra Alejandra, Weil Daniel, Colangelo Luigi.....	41
<b>N° 22 Upper eyelid retraction in Grave's disease</b>	
Magnano Paula, Billagra Alejandra, Avila Jorge, Racana Ignacio, Tula Romina, Weil Daniel .....	43
<b>N° 23 Ophthalmomyiasis and skin cancer</b>	
Billagra Alejandra, Matach Gustavo, Ortiz Basso Tomas .....	44
<b>N°24 Treatment of dry eye syndrome in young age glaucoma</b>	
Claudia Guarracino, Alessandra Trotta, Giuseppe Maisto .....	45
<b>N°25 Riboflavin-based staining of the epithelial ocular surface: a new diagnostic opportunity</b>	
Salvatore Troisi, Ciro Caruso, Elisabetta Chiariello Vecchio, Decio Capobianco, Paola Vele, Salvatore Adriano Cirillo .....	46
<b>N°26 The Schola Medica Salernitana and Ophthalmology</b>	
Giovanni Leo, Rossella Russo, Giusy Sbaglio, Attimonelli Pasquale, Luca Barile, Orazio Campagna .....	48
<b>N°27 The Aesthetic Medicine in the Schola Medica Salernitana: Trotula De Ruggiero</b>	
Giovanni Leo, Rossella Russo, Valeria Russo, Lorena Manzur, Maria Cristiana Abate, Catello Russo .....	50



## **PREFACE**

We are delighted to introduce this booklet of The IV International Course on Orbitoplasty and Neurophthalmology in Amalfi Coast, organized by Luigi Colangelo MD, our previous pupil in Chieti and Buenos Aires.

Flavio Gioia from Amalfi is attributed to define the magnetic compass finding the magnetic North and the Polaris in the Ursa minor constellation, whilst more or less in the same middle-age Vikings use the 'Iceland sunstone' [a calcite mineral, like the same that allowed trilobites develop the first simplest eye in the deep sea on Cambrian era] to find the sun behind the clouds, depolarizing light. The Schola Medica Salernitana, considered one of the oldest in modern civilization, together with Bologna and Paris, was empowered to release the licentia docendi that enabled the holder to teach ubique locorum ac per universum terrarum orbem. The aim was an holistic medicine in reference with the classical Galenic – Hippocratic tradition: the human body considered in its entirety as a microcosm within the macrocosm of nature, a tetradic holistic allopathic doctrine where contrariis contraria curantur. Astonishing for this age, the inner role of women who exercised their medical profession and wrote medicine treaties, as reported here inside.

The Amalfi coast is an unique landscape, loved by roman emperors, poets, musicians, dancers, artist from all over the world, where around the idea to confidentially share scientific experiences in a friendship atmosphere, he was able to connect surgeon and researchers from all over the world, deeply involved in these Ophthalmic sub-Specialities.

Everyone of these loves the Luigi Colangelo friendship, honouring his own study journey in Italy, Argentina, North America, Spain, North Europe and Portugal.

To reach the third biennial edition exposing the printed poster abstracts recollection to the scientific community, means the opportunity to maintain the memory of the exchanged knowledge together with Dermatologist and Plastic, Aesthetic, ENT and Maxillofacial Surgeons from one side and Neurophthalmologist and basic Researchers on the other, offering the opportunity to be quoted by the ISBN, as a respectful attitude to their studies, selected by the peer Scientific Committee.

The meeting was successful, as ever for his previous ones in the Amalfi coast. Luigi and the entire staff made a great personal effort to offer everyone the warmest taste of italian hospitality. The biennial invited lecture dedicated to his father Francesco Colangelo MD, gynaecologist and Maiori longstanding mayor, is the cherry on the cake as unforgettable flavour of this meeting. Pulchritudo animi bonum vinculum est inter homines.

**Pier Enrico Gallenga**

*FEBO*  
*G. D'Annunzio University Chieti*  
**ITALY**

**Daniel Weil**

*Jefe de la Sección de Orbita y Plástica Ocular del*  
*Servicio de Oftalmología. Hospital de Clínicas*  
*"José de San Martín" de la Universidad de Buenos Aires*  
**ARGENTINA**



## **‘FRANCESCO COLANGELO’ AWARD LECTURE**



As usual, this year too, I've organized an Award Lecture dedicated to my father, who from 1957 to 1997 practiced the medical profession on the Amalfi Coast with great passion.

**2007**

**EYE & POSTURE**



PIER ENRICO GALLENGA was born in Parma by Riccardo, ophthalmology professor, who held the chair of his grandfather Camillo, already Rector of the University.

He is author of more than 300 publications, 3 monographs and has a lot of participation to international congresses.

More than 30.000 operations realized in the structures of Chieti University where, from 1979, he manages the ophthalmological operating unit and the Ophthalmological Institute.

Deputy president of SOI 2003-2004 Chairman Ethics committee SOI from 2005

Medal of Strampelli lecture SOI 2005.

From 1995 he has founded and chairs the National Federation of Ethics Committees

**2009**

**SOME CONSIDERATION ON THE COMPARATIVE ANATOMY OF THE ORBIT**



DANIEL WEIL was born in Buenos Aires; he is teachers seconded as well as founder and director of the Orbital surgery section of the Department of Ophthalmology at the University of Buenos Aires. He is also teacher at the Sociedad Argentina de Oftalmologia, at the American Association of Ophthalmology, at Universidad del Salvador, at Maimonides University and at the Argentinian Catholic University.

He's honorary member of SOI and of the Brazilian Society of Ocular Plastic.

Past president of the Sociedad Argentina de Plastica Ocular as well as of the Sociedad Argentine de Oftalmologia.

He's member of the American Society of Ophthalmology and of the American Academy of Ophthalmology.

**2013**

## ORBITAL SURGERY: INDICATIONS & OPPORTUNITIES



LELIO BALDESCHI: MD PhD is "Chef de Clinique" and Clinical Professor in the Department of Ophthalmology, Catholic University of Louvain, at the Saint Luc Hospital of Bruxelles. His main field of interest is orbital and adnexal diseases. In such a field he trained residents in ophthalmology, nineteen international fellows, and had the opportunity to publish and lecturing at a wide variety of international congresses, and courses. He is member of SOI, ESOPRS, has been elected as honorary member of the APSOPRS, serves in the steering committee of the EUGOGO, is for the second time President of SICOP; he's the Secretary and the Elected President of the ISD&DE.

**2016**

## ANATOMY AND PHISIOPATHOLOGY OF LACRIMAL DUCTS



FAUSTO TRIVELLA: He was born in in Pisa in June 1957 , graduated in Medicine and Surgery in Pisa with honors, and specialized in Firenze in Ophthalmology with honors . During his post graduate work, he specializes in Eyelid orbit surgery as assistant in training in the Section of Craniofacial Surgery of Dr .Tessier at Hospital Foch of Paris, where he stayed for three years becoming a pupil of Doctor Krastinova with which he work also at Clinique Chateau de la Maye in Versailles. Assistant at the Teaching hospital of Ophthalmology in Florence, where he manage the Unit of Eyelid orbit surgery. Then he was moved to the Department of Ophthalmic surgery in Pisa where he manage the Unit of Oculoplastic surgery. Since 7 years he is head physician of Department of Ophthalmology of Lucca Hospital Since 20 years, he is consultant for neurosurgery and pediatric ophthalmology of Children hospital Meyer in Florence Winner of different prizes, among them SICOP award, the society of which he is also founder. Now he lives in Pisa and is engaged with Susanna with which he has had two children: Alessandro and Antonio.



# **EVALUATION OF THE RADIAL PERIPAPILLARY CAPILLARY NETWORK IN CONGENITAL OPTIC DISC ANOMALIES WITH OPTICAL COHERENCE TOMOGRAPHY ANGIOGRAPHY.**

Gilda Cennamo MD, Claudia Rossi MD, Pasquale Ruggiero MD and Giovanni Cennamo.

Eye Clinic, Department of Neurosciences, Reproductive Sciences and Dentistry, University of Naples Federico II, Naples, Italy

## **Goal**

To evaluate the radial peripapillary capillary network with optical coherence tomography angiography (angio-OCT) in Morning Glory Syndrome (MGS), optic disc colobomas and optic disc pits.

## **Materials and methods**

Fifteen eyes of 15 patients with congenital optic disc anomalies were enrolled in this study. All patients underwent optical coherence tomography angiography (angio-OCT). The scans were centered on optic discs.

## **Results**

The mean age at presentation was 33 years (range, 19–50 years). Congenital optic disc anomalies were identified in all 15 eyes. Three eyes had the characteristic fundoscopic signs of MGS, and angio-OCT scans of the peripapillary retina revealed a dense microvascular network. Optic disc colobomas were identified in 5 eyes, and the characteristic fundoscopic signs of optic pits in 7 eyes. Angio-OCT showed the absence of radial peripapillary microvascular network in these 12 eyes.

## **Conclusions**

Angio-OCT is a safe and rapid imaging technique that could shed light on the pathogenesis of rare diseases of the optic disc. In fact, our finding that angio-OCT scans confirmed the presence of a peripapillary microvascular network only in MGS cases, supports the hypothesis that a primary neuroectodermal abnormality and a secondary mesenchymal abnormality leads to MGS.

## **Bibliography**

- Evaluation of morning glory syndrome with spectral optical coherence tomography and ecography. Cennamo Gil, de Crecchio G, Iaccarino G, Forte R, Cennamo G *Ophthalmology* 2010;117:1269-73

## **LEBER HEREDITARY OPTIC NEUROPATHY: A CASE REPORT.**

Gianluca Baldi, Giovanni Leo, Maurizio Cioffi, Luca Barile, Assunta Rosaria Cascone, Alfonso D'Alessandro.

- Presidio Ospedaliero "Umberto I", Nocera Inferiore

### **Goal**

To describe a case of Leber optic neuropathy in a young patient.

### **Materials and methods**

Case report

### **Results**

A 40-year-old boy presented with painless vision loss in both eyes. On clinical examination, his visual acuity was 2/10 in OD and counting fingers at 40 cm OS. Both pupils reacted normally to light. The findings on slit-lamp examination were all unremarkable. On the other hand, funduscopy after pupillary dilation showed atrophic papillas in both eyes. Genetic analysis demonstrated a point mutation of the mitochondrial gene ND4 (m11778G>A). He had no visual recovery after treatment with oral coenzyme Q10, vitamin B1, and citicoline.

### **Conclusions**

Leber hereditary optic neuropathy is one of the most common inherited optic neuropathies causing bilateral central vision loss. Molecular examination of mtDNA mutation can confirm the diagnosis of Leber neuropathy in clinically controversial patients.

### **Bibliography**

- Rasool N, Lessell S, Cestari DM. Leber Hereditary Optic Neuropathy: Bringing the Lab to the Clinic. *Semin Ophthalmol.* 2016; 31: 107-116.
- Newman NJ. Hereditary optic neuropathies: from the mitochondria to the optic nerve. *Am J Ophthalmol* 2005; 140: 517-523.

## **FIRST ITALIAN FAMILY WITH AUTOSOMAL RECESSIVE RETINAL DYSTROPHY CAUSED BY MERTK MUTATION (RP38).**

Gallenga CE<sub>1</sub>, Lanciotti S<sub>2</sub>, Lattanzio D<sub>3</sub>, Bolletta E<sub>1</sub>, Parmeggiani F<sub>1</sub>, Perri P<sub>1</sub>,  
Giovannini A<sub>4</sub>, Novelli G<sub>2</sub>, Brancati F<sub>5</sub>.

1. Dept Biomedical Surgical Specialistic Sciences, section Ophthalmology, University of Ferrara
2. Dept Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy
3. Dept Exper Surgical Sciences, section Ophthalmology, G. D'Annunzio University, Chieti-Pescara
4. Dept Neurosciences, section Ophthalmology, Marche Polytechnic University, Ancona
5. Dept of Life, Health & Environmental Sciences, University of L'Aquila, L'Aquila, Italy

### **Goal**

To describe the genetic and ocular findings in an Italian 12-year-old girl showing cone-rod "pisciform-like" macular dystrophy and to broaden the molecular spectrum of *MERTK* gene

### **Materials and Methods**

A complete ophthalmological examination was carried out, implemented with: OCT, autofluorescence, FA, ICGA, electrophysiology (flicker- and pattern-ERG-, PEV), contrast sensitivity, chromatic sense, visual field (120p, 30/2), orthoptic evaluation. The patient was followed in different ophthalmological centers starting from age 4 and the final differential diagnosis between Stargardt disease and cone-rod dystrophy was pointed out at age of 12 and then collaboratively investigated by geneticist.

A panel of nearly 100 retinal dystrophy disease-causative genes was analyzed by next-generation sequencing techniques for diagnostic purposes. After a putative mutation was identified, Sanger sequencing confirmation of the distinct genomic fragment as well as segregation analysis in the family were performed. The pathogenic role of the mutation was predicted by *in silico* models.

### **Results**

The patient underwent craniotomy at age seven months for meningocele. Visual impairment was firstly noticed at age 6 years and progressively increase until stabilization at 2/10 bilaterally. Severe dysfunction of both photoreceptor systems involved the macula at an early stage. Other findings include: nictalopia, photoaversion, dyschromatopsia, visual field centrocecal mixed scotoma to the mid periphery, ERG b wave reduced in RE and absent in LE, reduced macular thickness (168  $\mu$ m) with RPE drusen-like changes and photoreceptors thinning with high

reflectance bodies below the outer limiting membrane, without CME, discrete dot-like autofluorescent deposits hypofluorescent on ICGA (hallmark of MERTK-specific retinal dystrophies). A nucleotide change affecting an obligatory splice site of intron 10 of the *MERTK* gene (c.1604+5G>A) was depicted at the homozygote state in the proband, while her parents were heterozygote carriers.

### Conclusions

We report a novel homozygous splicing mutation in the *MERTK* gene (OMIM\*604705) and emphasize distinct ophthalmologic features of this rare cause of autosomal recessive retinal dystrophy (overall around 1%). In the era of personalized medicine, a multidisciplinary approach is warranted to reach a precise diagnosis of retinal dystrophy and plan gene-based pharmacological trials.

### Bibliography

- Gal A, Li Y, Thompson DA, Weir J, Orth U, Jacobson SG, Apfelstedt-Sylla E, Vollrath D. Mutation in MERTK, the human orthologue of the RCS rat retinal dystrophy gene, cause retinitis pigmentosa. *Nature Genet.* 2000;26: 270-271.
- Parinot C, Nandrot EF. A comprehensive review of mutations in the MERTK protooncogene. *Adv Exp Med Biol.* 2016;854:259-65.
- Jinda W, Pongvarin N, Taylor TD, Suzuki Y, Thongnoppakhun W, Limwongse C, Lertrit P, Suriyaphol P, Atchaneeyasakul LO. A novel start codon mutation of the MERTK gene in a patient with retinitis pigmentosa. *Molecular Vision.* 2016;22:342-51.
- Abigail T Fahim, Stephen P Daiger, and Richard G Weleber. Retinitis Pigmentosa Overview. *GeneReviews®.* March 21, 2013.
- Brea-Fernández AJ, Pomares E, Brión MJ, Marfany G, Blanco MJ, Sánchez-Salorio M, González-Duarte R, Carracedo A. Novel splice donor site mutation in MERTK gene associated with retinitis pigmentosa. *Br J Ophthalmol.* 2008;92(10):1419-23.
- Mackay DS, Henderson RH, Sergouniotis PI, Li Z, Moradi P, Holder GE, Waseem N, Bhattacharya SS, Aldahmesh MA, Alkuraya FS, Meyer B, Webster AR, Moore AT. Novel mutations in MERTK associated with childhood onset rod-cone dystrophy. *Molecular Vision.* 2010;16:369-77.
- Ksantini M, Lafont E, Bocquet B, Meunier I, Hamel CP. Homozygous mutation in MERTK causes severe autosomal recessive retinitis pigmentosa. *Eur J Ophthalmol.* 2012;22(4):647-53.
- Charbel Issa P, Bolz HJ, Ebermann I, Domeier E, Holz FG, Scholl HP. Characterization of severe rod-cone dystrophy in a consanguineous family with a splice site mutation in the MERTK gene. *British journal of ophthalmology* 2009;93(7):920-5.
- Ghazi NG, Abboud EB, Nowilaty SR, Alkuraya H, Alhommadi A, Cai H, Hou R, Deng WT, Boye SL, Almaghamsi A, Al Saikhan F, Al-Dhibi H, Birch D, Chung C, Colak D,

LaVail MM, Vollrath D, Erger K, Wang W, Conlon T, Zhang K, Hauswirth W, Alkuraya FS. Treatment of retinitis pigmentosa due to MERTK mutations by ocular subretinal injection of adeno-associated virus gene vector: results of a phase I trial. *Hum Genet.* 2016 Mar;135(3):327-43.



## **IMPROVING VA USING GABOR PATCH TECHNOLOGY IN STARGARDT DISEASE: A RETROSPECTIVE STUDY**

Mincarelli C<sup>°</sup>, Gallenga CE\*, Colangelo L<sup>”</sup>, Lobefalo L<sup>^</sup>, Rapinese M<sup>^</sup>, Di Carlo PL<sup>^</sup>, Nuri M<sup>^</sup>, Perri P\*, Parmeggiani F\*.

<sup>°</sup> CNO Orthoptic service, Pescara, Italy

\* Dept Biomedical Surgical Specialistic Sciences, section Ophthalmology, University of Ferrara, Italy

" Salus Health Center, section Ophthalmology, Battipaglia, Italy

<sup>^</sup> Tripoli Eye Hospital, Tripoli, Libia

<sup>^</sup> Dept Exper Surgical Sciences, section Ophthalmology, G D'Annunzio University, Chieti-Pescara, Italy

### **Goal**

To find the efficacy of a non-invasive, patient-specific treatment based on visual stimulation and facilitation of neural connections responsible for vision treatment in enhancing Best Corrected Visual Acuity (BCVA) in patients with Stargardt disease. As already demonstrated in other applications such as adult amblyopia, the technology involves the use of an internet-based computer generated visual training exercise regime using sets of patient-specific stimuli based on Gabor patches, to improve contrast sensitivity and, therefore, visual acuity.

### **Materials and Methods**

5 Stargardt patients (10 eyes) verified at Ophthalmoscopy/ERG/ FA/ICG/OCT/Ishihara tables, 3 male, 2 female, mean aged 35 (+- 11.11 y,c), caucasians, were selected and informed according to the Helsinki declaration; exclusion criteria were age (<18 > 50ny. o.), other ocular disease impairing the visual axis, neurological disease or mental impairment. Inclusion criteria: BCVA in both eyes stable in the last 6 month. Gabor patch technology involves the use of an internet-based computer generated visual training exercise regime using sets of patient-specific stimuli based on Gabor patches, to sharpen contrast sensitivity and visual acuity. The fundamental stimulation-control technique is called "Lateral Masking", where collinearly oriented flanking Gabors are displayed in addition to the target Gabor image. This NeuroVision Treatment System is a software-based, interactive system tailored and continuously adaptive to the individual visual abilities. In the first stage, the subject is exposed to a set of visual perception tasks, aimed to analyze and identify each subject's neural inefficiencies or deficiencies. Based on this analysis, a treatment plan is initialized, and subject's V1 cortical area receives patients-specific stimuli in a controlled environment.

### **Results**

Mean BCVA, for the 10 eyes, has raised from 0.81 (+- 0.17) logMar to 0.49 (+-0.07) logMar. Mean ETDRS' chart lines improvement has been 3.18 +-2.05.

One subject has quit the treatment for pregnancy.

## **Conclusions**

Results suggest that Gabor patch treatment improves BCVA in Stargard patients. This improvement appears to be retained for at least 12 months after treatment, compatibly with the disease itself progression.

## **Bibliography**

- Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol. (Lond.)* 160, 106-154 (1962).
- Gallenga PE, Rapinese M, Lobefalo L, Cerulli L, Cedrone C. Low vision and blindness in Abruzzo. *New Trends Ophthalmol* 5,137-142 (1989).
- Rapinese M, Lobefalo L, Marsilio T, D'Aurelio AP, Lapi G. Influenza dei fattori oggettivi e soggettivi nella riabilitazione del paziente ipovedente. *Minerva Oftalmologica* 36, 149-52 (1994).
- Rapinese M, Lobefalo L, Baiocchi A, Crudeli CF, Di Carlo PL, Di Crescenzo F, Marsilio T, Mastropasqua L, Gallenga PE. Biofeedback therapy in ambliopic adults. *Ophth Res* 30 S1 (1998).
- Polat, U., Mizobe, K., Pettet, M. W., Kasamatsu, T. & Norcia, A. M. Collinear stimuli regulate visual responses depending on cell's contrast threshold. *Nature* 391, 580-4 (1998).
- Polat, U., Ma-Naim, T. Belkin, M. Sagi, D. Improving vision in adult amblyopia by perceptual learning. *PNAS* 101, 6692-97 (2004).

# **EXENTERATION OF ORBIT IN A TERTIARY EYE HOSPITAL - INDICATION AND OUTCOMES**

(1,2) Silvana A. Schellini, (1,3) Alicia Galindo-Ferreiro, (1) Rajiv Khandekar, (4) Patricia M. Akaishi, (4) Antonio Augusto V. Cruz, (5) Yereña Muiños

<sup>1</sup> KING KHALED EYE SPECIALIST HOSPITAL, RIYADH, SAUDI ARABIA;

<sup>2</sup> FACULDADE DE MEDICINA DE BOTUCATU - UNESP, SÃO PAULO, BRAZIL;

<sup>3</sup> COMPLEJO HOSPITALARIO PALENCAL, SPAIN

<sup>4</sup> FACULDADE DE MEDICINA DE RIBEIRÃO PRETO – USP, SAO PAULO, BRAZIL

<sup>5</sup> HOSPITAL NUESTRA SEÑORA DE FÁTIMA. VIGO, SPAIN

## **Goal**

To present indications, type of surgery and outcomes of orbital exenteration (OE) at a tertiary eye hospital in 20 years.

## **Materials and methods**

This was a retrospective review of health records involving all patients who underwent to OE between 1994 and 2014 at KKESH. Demographic data, ocular status prior to surgery, details of exenteration, histopathological diagnosis, adjunctive treatment, follow-up examination findings, recurrences, complications and their management were noted.

## **Results**

Orbital exenteration was performed on 60 orbits of 60 patients. Median age 68.8 25% quartile 57.5 minimum 2 years Maximum 96. The indications for OE were malignancies in 59(98.3%) cases. The commonest malignancies noted in this series were SCC (60%, 36/60) and BCC eyelids with an orbital involvement (15%, 9/60). A total OE was required for 17 patients, a subtotal for further 41 and two patients had extended OE.

## **Conclusions**

OE is usually indicated to treat malignant tumors and SCC was the single most common neoplasm responsible for OE at KKESH.

## **Bibliography**

1. Pollack RS. Orbital and periorbital tumors. Indications for exenteration. *California medicine*. 1959;91:207-11.
2. Nemet AY, Martin P, Bengner R, Kourt G, Sharma V, Ghabrial R, et al. Orbital exenteration: a 15-year study of 38 cases. *Ophthalmic plastic and reconstructive surgery*. 2007;23(6):468-72.
3. Tyers AG. Orbital exenteration for invasive skin tumours. *Eye*. 2006;20(10):1165-70.
4. Pushker N, Kashyap S, Balasubramanya R, Bajaj MS, Sen S, Betharia SM, et al. Pattern of orbital exenteration in a tertiary eye care centre in India. *Clinical & experimental ophthalmology*. 2004;32(1):51-4.

5. Bartley GB, Garrity JA, Waller RR, Henderson JW, Ilstrup DM. Orbital exenteration at the Mayo Clinic. 1967-1986. *Ophthalmology*. 1989;96(4):468-73.
6. Goldberg RA, Kim JW, Shorr N. Orbital exenteration: results of an individualized approach. *Ophthalmic plastic and reconstructive surgery*. 2003;19(3):229-36.
7. Shields JA, Shields CL, Demirci H, Honavar SG, Singh AD. Experience with eyelid-sparing orbital exenteration: the 2000 Tullos O. Coston Lecture. *Ophthalmic plastic and reconstructive surgery*. 2001;17(5):355-61.
8. Looi A, Kazim M, Cortes M, Rootman J. Orbital reconstruction after eyelid- and conjunctiva-sparing orbital exenteration. *Ophthalmic plastic and reconstructive surgery*. 2006;22(1):1-6.
9. Yeatts RP. The esthetics of orbital exenteration. *American journal of ophthalmology*. 2005;139(1):152-3.
10. Rodrigues ML, Kohler HF, Faria JC, Ikeda MK, Vartanian JG, Kowalski LP. Reconstruction after extended orbital exenteration using a fronto-lateral flap. *International journal of oral and maxillofacial surgery*. 2009;38(8):850-4.
11. Suarez C, Ferlito A, Lund VJ, Silver CE, Fagan JJ, Rodrigo JP, et al. Management of the orbit in malignant sinonasal tumors. *Head & neck*. 2008;30(2):242-50.
12. Croce A, Moretti A, D'Agostino L, Zingariello P. Orbital exenteration in elderly patients: personal experience. *Acta otorhinolaryngologica Italica : organo ufficiale della Societa italiana di otorinolaringologia e chirurgia cervico-facciale*. 2008;28(4):193-9.
13. Ackuaku-Dogbe E. Review of orbital exenterations in Korle-Bu teaching hospital. *Ghana medical journal*. 2011;45(2):45-9.
14. Frunza A, Slavescu D, Zamfirescu D, Stanculescu L, Grintescu I, Enache V, et al. Orbital exenteration - a salvage procedure? *Romanian journal of morphology and embryology = Revue roumaine de morphologie et embryologie*. 2013;54(4):1161-7.
15. Hoffman GR, Jefferson ND, Reid CB, Eisenberg RL. Orbital Exenteration to Manage Infiltrative Sinonasal, Orbital Adnexal, and Cutaneous Malignancies Provides Acceptable Survival Outcomes: An Institutional Review, Literature Review, and Meta-Analysis. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2015.
16. Kaur A, Khattri M, Jaiswal V. Pattern of orbital exenteration at a tertiary eye care centre in North India. *Nepalese journal of ophthalmology : a biannual peer-reviewed academic journal of the Nepal Ophthalmic Society : NEPJOPH*. 2012;4(1):64-7.
17. Maheshwari R. Review of orbital exenteration from an eye care centre in Western India. *Orbit*. 2010;29(1):35-8.
18. Nassab RS, Thomas SS, Murray D. Orbital exenteration for advanced periorbital skin cancers: 20 years experience. *Journal of plastic, reconstructive & aesthetic surgery : JPRAS*. 2007;60(10):1103-9.
19. Qassemayar A, Aljudaibi N, Wavreille O, Mortier L, Martinot-Duquennoy V, Guerreschi P. Orbital exenteration and periorbital skin cancers. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2014;72(4):811-6.
20. Soysal HG. Orbital exenteration: a 10-year experience of a general oncology hospital. *Orbit*. 2010;29(3):135-9.
21. Ben Simon GJ, Schwarcz RM, Douglas R, Fiaschetti D, McCann JD, Goldberg RA. Orbital exenteration: one size does not fit all. *American journal of ophthalmology*. 2005;139(1):11-7.

22. Rahman I, Cook AE, Leatherbarrow B. Orbital exenteration: a 13 year Manchester experience. *The British journal of ophthalmology*. 2005;89(10):1335-40.
23. Ali MJ, Pujari A, Dave TV, Kaliki S, Naik MN. Clinicopathological profile of orbital exenteration: 14 years of experience from a tertiary eye care center in South India. *International ophthalmology*. 2015.
24. Atique-Tacla M, Paves L, Pereira MD, Manso PG. [Exenteration: a retrospective study]. *Arquivos brasileiros de oftalmologia*. 2006;69(5):679-82. Exenteracao: estudo retrospectivo.
25. Levin PS, Dutton JJ. A 20-year series of orbital exenteration. *American journal of ophthalmology*. 1991;112(5):496-501.
26. Mohr C, Esser J. Orbital exenteration: surgical and reconstructive strategies. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 1997;235(5):288-95.
27. Naquin HA. Exenteration of the orbit. *AMA archives of ophthalmology*. 1954;51(6):850-62.
28. Rathbun JE, Beard C, Quickert MH. Evaluation of 48 cases of orbital exenteration. *American journal of ophthalmology*. 1971;72(1):191-9.
29. Simons JN, Robinson DW, Masters FW. Malignant tumors of the orbit and periorbital structures treated by exenteration. *Plastic and reconstructive surgery*. 1966;37(2):100-4.
30. Putterman AM. Orbital exenteration with spontaneous granulation. *Archives of ophthalmology*. 1986;104(1):139-40.
31. Levin PS, Ellis DS, Stewart WB, Toth BA. Orbital exenteration. The reconstructive ladder. *Ophthalmic plastic and reconstructive surgery*. 1991;7(2):84-92.
32. Reese AB. Exenteration of the orbit; with transplantation of the temporalis muscle. *American journal of ophthalmology*. 1958;45(3):386-90.
33. Reese AB, Jones IS. Exenteration of the orbit and repair by transplantation of the temporalis muscle. *American journal of ophthalmology*. 1961;51:217-27.
34. Dortzbach RK, Hawes MJ. Midline forehead flap in reconstructive procedures of the eyelids and exenterated socket. *Ophthalmic surgery*. 1981;12(4):257-68.
35. Shore JW, Burks R, Leone CR, Jr., McCord CD, Jr. Dermis-fat graft for orbital reconstruction after subtotal exenteration. *American journal of ophthalmology*. 1986;102(2):228-36.
36. Mauriello JA, Jr., Han KH, Wolfe R. Use of autogenous split-thickness dermal graft for reconstruction of the lining of the exenterated orbit. *American journal of ophthalmology*. 1985;100(3):465-7.
37. Shields JA, Shields CL, Suvarnamani C, Tantisira M, Shah P. Orbital exenteration with eyelid sparing: indications, technique, and results. *Ophthalmic surgery*. 1991;22(5):292-7.

# **EXPANSIVE HYDROGEL PELLETS IN THE TREATMENT OF POST ENUCLEATION/EVSCERATION SOCKET SYNDROME**

Delmas Juliette, Lossouran Adrien, Adenis Jean-Paul, Robert Pierre-Yves

1. Centre Hospitalier Universitaire Dupuytren – Limoges - France

## **Goal**

To evaluate the post-operative result after orbital injection of expansive hydrogel pellets, in the treatment of enophthalmos secondary to PESS.

## **Materials and methods**

This study was carried out between January 2010 and September 2015, in the University Hospital Dupuytren Limoges – France. Twenty surgeries were performed in 16 patients (20 procedures). Intra-orbital injection of the hydrogel pellets was performed under general anesthesia, behind the hydroxyapatite orbital implant, through small conjunctival incisions (1.5 mm). The volume of a single hydrogel pellet volume is 0.024 ml, and it increases tenfold after intra-orbital rehydration, after 24 to 48 hours.

## **Results**

The average enophthalmos rose from 13.35 +/- 3.69 (6-19) mm pre-injection to 14.98 +/- 3.69 (7-20) post-injection. The average number of injected pellets was 7.45 +/- 2.76 (3-15) per procedure. Three sockets (18.75%) needed a second surgery, with the same procedure. Esthetic result after fitting a new prosthetic device was satisfactory in 100% of the cases. Half of the patients felt a post operative pain, successfully treated by analgesic. One of our patient presented a sub-cutaneous implant migration, although not requiring surgical correction.

## **Conclusions**

This procedure appears sure, safe, quick, without risk of any serious complication, and leading to good results. It can be performed on ambulatory surgery. The expected post-operative increase of enophthalmos is 1 to 3 mm. Up to 10 or 11 pellets can be injected but more may lead to severe post-operative pain. Long term results seem to stable over time but a longer follow-up is necessary to evaluate the risk of late implant migration or extrusion.

## **Bibliography**

- Tyers AG, Collin JR. Orbital implants and post enucleation socket syndrome. *Trans Ophthalmol Soc U K.* 1982 Apr;102 (Pt 1):90–2.
- Wiese KG, Vogel M, Guthoff R, Gundlach KK. Treatment of congenital anophthalmos with self-inflating polymer expanders: a new method. *J Cranio-Maxillo-fac Surg Off Publ Eur Assoc Cranio-Maxillo-fac Surg.* 1999 Apr;27(2):72–6.
- Gundlach KKH, Guthoff RF, Hingst VHM, Schittkowski MP, Bier UC. Expansion of the socket and orbit for congenital clinical anophthalmia. *Plast Reconstr Surg.* 2005 Oct;116(5):1214–22.

## **LEBER HEREDITARY OPTIC NEUROPATHY: A CASE REPORT.**

Gianluca Baldi, Giovanni Leo, Maurizio Cioffi, Luca Barile, Assunta Rosaria Cascone, Alfonso D'Alessandro.

- Presidio Ospedaliero "Umberto I", Nocera Inferiore

### **Goal**

To describe a case of Leber optic neuropathy in a young patient.

### **Materials and methods**

Case report

### **Results**

A 40-year-old boy presented with painless vision loss in both eyes. On clinical examination, his visual acuity was 2/10 in OD and counting fingers at 40 cm OS. Both pupils reacted normally to light. The findings on slit-lamp examination were all unremarkable. On the other hand, funduscopy after pupillary dilation showed atrophic papillas in both eyes. Genetic analysis demonstrated a point mutation of the mitochondrial gene ND4 (m11778G>A). He had no visual recovery after treatment with oral coenzyme Q10, vitamin B1, and citicoline.

### **Conclusions**

Leber hereditary optic neuropathy is one of the most common inherited optic neuropathies causing bilateral central vision loss. Molecular examination of mtDNA mutation can confirm the diagnosis of Leber neuropathy in clinically controversial patients.

### **Bibliography**

- Rasool N, Lessell S, Cestari DM. Leber Hereditary Optic Neuropathy: Bringing the Lab to the Clinic. *Semin Ophthalmol.* 2016; 31: 107-116.
- Newman NJ. Hereditary optic neuropathies: from the mitochondria to the optic nerve. *Am J Ophthalmol* 2005; 140: 517-523.

## **FIRST ITALIAN FAMILY WITH AUTOSOMAL RECESSIVE RETINAL DYSTROPHY CAUSED BY MERTK MUTATION (RP38).**

Gallenga CE<sub>1</sub>, Lanciotti S<sub>2</sub>, Lattanzio D<sub>3</sub>, Bolletta E<sub>1</sub>, Parmeggiani F<sub>1</sub>, Perri P<sub>1</sub>,  
Giovannini A<sub>4</sub>, Novelli G<sub>2</sub>, Brancati F<sub>5</sub>.

1. Dept Biomedical Surgical Specialistic Sciences, section Ophthalmology, University of Ferrara
2. Dept Biomedicine and Prevention, University of Rome Tor Vergata, Rome, Italy
3. Dept Exper Surgical Sciences, section Ophthalmology, G. D'Annunzio University, Chieti-Pescara
4. Dept Neurosciences, section Ophthalmology, Marche Polytechnic University, Ancona
5. Dept of Life, Health & Environmental Sciences, University of L'Aquila, L'Aquila, Italy

### **Goal**

To describe the genetic and ocular findings in an Italian 12-year-old girl showing cone-rod "pisciform-like" macular dystrophy and to broaden the molecular spectrum of *MERTK* gene

### **Materials and Methods**

A complete ophthalmological examination was carried out, implemented with: OCT, autofluorescence, FA, ICGA, electrophysiology (flicker- and pattern-ERG-, PEV), contrast sensitivity, chromatic sense, visual field (120p, 30/2), orthoptic evaluation. The patient was followed in different ophthalmological centers starting from age 4 and the final differential diagnosis between Stargardt disease and cone-rod dystrophy was pointed out at age of 12 and then collaboratively investigated by geneticist.

A panel of nearly 100 retinal dystrophy disease-causative genes was analyzed by next-generation sequencing techniques for diagnostic purposes. After a putative mutation was identified, Sanger sequencing confirmation of the distinct genomic fragment as well as segregation analysis in the family were performed. The pathogenic role of the mutation was predicted by *in silico* models.

### **Results**

The patient underwent craniotomy at age seven months for meningocele. Visual impairment was firstly noticed at age 6 years and progressively increase until stabilization at 2/10 bilaterally. Severe dysfunction of both photoreceptor systems involved the macula at an early stage. Other findings include: nictalopia, photoaversion, dyschromatopsia, visual field centrocecal mixed scotoma to the mid periphery, ERG b wave reduced in RE and absent in LE, reduced macular thickness (168  $\mu$ m) with RPE drusen-like changes and photoreceptors thinning with high



reflectance bodies below the outer limiting membrane, without CME, discrete dot-like autofluorescent deposits hypofluorescent on ICGA (hallmark of MERTK-specific retinal dystrophies). A nucleotide change affecting an obligatory splice site of intron 10 of the *MERTK* gene (c.1604+5G>A) was depicted at the homozygote state in the proband, while her parents were heterozygote carriers.

### **Conclusions**

We report a novel homozygous splicing mutation in the *MERTK* gene (OMIM\*604705) and emphasize distinct ophthalmologic features of this rare cause of autosomal recessive retinal dystrophy (overall around 1%). In the era of personalized medicine, a multidisciplinary approach is warranted to reach a precise diagnosis of retinal dystrophy and plan gene-based pharmacological trials.

### **Bibliography**

- Gal A, Li Y, Thompson DA, Weir J, Orth U, Jacobson SG, Apfelstedt-Sylla E, Vollrath D. Mutation in MERTK, the human orthologue of the RCS rat retinal dystrophy gene, cause retinitis pigmentosa. *Nature Genet.* 2000;26: 270-271.
- Parinot C, Nandrot EF. A comprehensive review of mutations in the MERTK protooncogene. *Adv Exp Med Biol.* 2016;854:259-65.
- Jinda W, Pongvarin N, Taylor TD, Suzuki Y, Thongnoppakhun W, Limwongse C, Lertrit P, Suriyaphol P, Atchaneeyasakul LO. A novel start codon mutation of the MERTK gene in a patient with retinitis pigmentosa. *Molecular Vision.* 2016;22:342-51.
- Abigail T Fahim, Stephen P Daiger, and Richard G Weleber. Retinitis Pigmentosa Overview. *GeneReviews®.* March 21, 2013.
- Brea-Fernández AJ, Pomares E, Brión MJ, Marfany G, Blanco MJ, Sánchez-Salorio M, González-Duarte R, Carracedo A. Novel splice donor site mutation in MERTK gene associated with retinitis pigmentosa. *Br J Ophthalmol.* 2008;92(10):1419-23.
- Mackay DS, Henderson RH, Sergouniotis PI, Li Z, Moradi P, Holder GE, Waseem N, Bhattacharya SS, Aldahmesh MA, Alkuraya FS, Meyer B, Webster AR, Moore AT. Novel mutations in MERTK associated with childhood onset rod-cone dystrophy. *Molecular Vision.* 2010;16:369-77.
- Ksantini M, Lafont E, Bocquet B, Meunier I, Hamel CP. Homozygous mutation in MERTK causes severe autosomal recessive retinitis pigmentosa. *Eur J Ophthalmol.* 2012;22(4):647-53.
- Charbel Issa P, Bolz HJ, Ebermann I, Domeier E, Holz FG, Scholl HP. Characterization of severe rod-cone dystrophy in a consanguineous family with a splice site mutation in the MERTK gene. *British journal of ophthalmology* 2009;93(7):920-5.
- Ghazi NG, Abboud EB, Nowilaty SR, Alkuraya H, Alhommadi A, Cai H, Hou R, Deng WT, Boye SL, Almaghamsi A, Al Saikhan F, Al-Dhibi H, Birch D, Chung C, Colak D,

LaVail MM, Vollrath D, Erger K, Wang W, Conlon T, Zhang K, Hauswirth W, Alkuraya FS. Treatment of retinitis pigmentosa due to MERTK mutations by ocular subretinal injection of adeno-associated virus gene vector: results of a phase I trial. *Hum Genet.* 2016 Mar;135(3):327-43.

## **IMPROVING VA USING GABOR PATCH TECHNOLOGY IN STARGARDT DISEASE: A RETROSPECTIVE STUDY**

Mincarelli C<sup>°</sup>, Gallenga CE\*, Colangelo L<sup>”</sup>, Lobefalo L<sup>^</sup>, Rapinese M<sup>^</sup>, Di Carlo PL<sup>^</sup>, Nuri M<sup>^</sup>, Perri P\*, Parmeggiani F\*.

<sup>°</sup> CNO Orthoptic service, Pescara, Italy

\* Dept Biomedical Surgical Specialistic Sciences, section Ophthalmology, University of Ferrara, Italy

" Salus Health Center, section Ophthalmology, Battipaglia, Italy

<sup>^</sup> Tripoli Eye Hospital, Tripoli, Libia

<sup>^</sup> Dept Exper Surgical Sciences, section Ophthalmology, G D'Annunzio University, Chieti-Pescara, Italy

### **Goal**

To find the efficacy of a non-invasive, patient-specific treatment based on visual stimulation and facilitation of neural connections responsible for vision treatment in enhancing Best Corrected Visual Acuity (BCVA) in patients with Stargardt disease. As already demonstrated in other applications such as adult amblyopia, the technology involves the use of an internet-based computer generated visual training exercise regime using sets of patient-specific stimuli based on Gabor patches, to improve contrast sensitivity and, therefore, visual acuity.

### **Materials and Methods**

5 Stargardt patients (10 eyes) verified at Ophthalmoscopy/ERG/ FA/ICG/OCT/Ishihara tables, 3 male, 2 female, mean aged 35 (+- 11.11 y,c), caucasians, were selected and informed according to the Helsinki declaration; exclusion criteria were age (<18 > 50ny. o.), other ocular disease impairing the visual axis, neurological disease or mental impairment. Inclusion criteria: BCVA in both eyes stable in the last 6 month. Gabor patch technology involves the use of an internet-based computer generated visual training exercise regime using sets of patient-specific stimuli based on Gabor patches, to sharpen contrast sensitivity and visual acuity. The fundamental stimulation-control technique is called “Lateral Masking”, where collinearly oriented flanking Gabors are displayed in addition to the target Gabor image. This NeuroVision Treatment System is a software-based, interactive system tailored and continuously adaptive to the individual visual abilities. In the first stage, the subject is exposed to a set of visual perception tasks, aimed to analyze and identify each subject’s neural inefficiencies or deficiencies. Based on this analysis, a treatment plan is initialized, and subject’s V1 cortical area receives patients-specific stimuli in a controlled environment.

### **Results**

Mean BCVA, for the 10 eyes, has raised from 0.81 (+- 0.17) logMar to 0.49 (+-0.07) logMar. Mean ETDRS’ chart lines improvement has been 3.18 +-2.05. One subject has quit the treatment for pregnancy.

## **Conclusions**

Results suggest that Gabor patch treatment improves BCVA in Stargard patients. This improvement appears to be retained for at least 12 months after treatment, compatibly with the disease itself progression.

## **Bibliography**

- Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. *J. Physiol. (Lond.)* 160, 106-154 (1962).
- Gallenga PE, Rapinese M, Lobefalo L, Cerulli L, Cedrone C. Low vision and blindness in Abruzzo. *New Trends Ophthalmol* 5,137-142 (1989).
- Rapinese M, Lobefalo L, Marsilio T, D'Aurelio AP, Lapi G. Influenza dei fattori oggettivi e soggettivi nella riabilitazione del paziente ipovedente. *Minerva Oftalmologica* 36, 149-52 (1994).
- Rapinese M, Lobefalo L, Baiocchi A, Crudeli CF, Di Carlo PL, Di Crescenzo F, Marsilio T, Mastropasqua L, Gallenga PE. Biofeedback therapy in ambliopic adults. *Ophth Res* 30 S1 (1998).
- Polat, U., Mizobe, K., Pettet, M. W., Kasamatsu, T. & Norcia, A. M. Collinear stimuli regulate visual responses depending on cell's contrast threshold. *Nature* 391, 580-4 (1998).
- Polat, U., Ma-Naim, T. Belkin, M. Sagi, D. Improving vision in adult amblyopia by perceptual learning. *PNAS* 101, 6692-97 (2004).

# EXENTERATION OF ORBIT IN A TERTIARY EYE HOSPITAL - INDICATION AND OUTCOMES

(1,2) Silvana A. Schellini, (1,3) Alicia Galindo-Ferreiro, (1) Rajiv Khandekar, (4) Patricia M. Akaishi, (4) Antonio Augusto V. Cruz, (5) Yerena Muiños

<sup>1</sup> KING KHALED EYE SPECIALIST HOSPITAL, RIYADH, SAUDI ARABIA;

<sup>2</sup> FACULDADE DE MEDICINA DE BOTUCATU - UNESP, SÃO PAULO, BRAZIL;

<sup>3</sup> COMPLEJO HOSPITALARIO PALENCAL, SPAIN

<sup>4</sup> FACULDADE DE MEDICINA DE RIBEIRÃO PRETO – USP, SAO PAULO, BRAZIL

<sup>5</sup> HOSPITAL NUESTRA SEÑORA DE FÁTIMA. VIGO, SPAIN

## Goal

To present indications, type of surgery and outcomes of orbital exenteration (OE) at a tertiary eye hospital in 20 years.

## Materials and methods

This was a retrospective review of health records involving all patients who underwent to OE between 1994 and 2014 at KKESH. Demographic data, ocular status prior to surgery, details of exenteration, histopathological diagnosis, adjunctive treatment, follow-up examination findings, recurrences, complications and their management were noted.

## Results

Orbital exenteration was performed on 60 orbits of 60 patients. Median age 68.8 25% quartile 57.5 minimum 2 years Maximum 96. The indications for OE were malignancies in 59(98.3%) cases. The commonest malignancies noted in this series were SCC (60%, 36/60) and BCC eyelids with an orbital involvement (15%, 9/60). A total OE was required for 17 patients, a subtotal for further 41 and two patients had extended OE.

## Conclusions

OE is usually indicated to treat malignant tumors and SCC was the single most common neoplasm responsible for OE at KKESH.

## Bibliography

1. Pollack RS. Orbital and periorbital tumors. Indications for exenteration. California medicine. 1959;91:207-11.
2. Nemet AY, Martin P, Bengner R, Kourt G, Sharma V, Ghabrial R, et al. Orbital exenteration: a 15-year study of 38 cases. Ophthalmic plastic and reconstructive surgery. 2007;23(6):468-72.
3. Tyers AG. Orbital exenteration for invasive skin tumours. Eye. 2006;20(10):1165-70.
4. Pushker N, Kashyap S, Balasubramanya R, Bajaj MS, Sen S, Betharia SM, et al. Pattern of orbital exenteration in a tertiary eye care centre in India. Clinical & experimental ophthalmology. 2004;32(1):51-4.

5. Bartley GB, Garrity JA, Waller RR, Henderson JW, Ilstrup DM. Orbital exenteration at the Mayo Clinic. 1967-1986. *Ophthalmology*. 1989;96(4):468-73.
6. Goldberg RA, Kim JW, Shorr N. Orbital exenteration: results of an individualized approach. *Ophthalmic plastic and reconstructive surgery*. 2003;19(3):229-36.
7. Shields JA, Shields CL, Demirci H, Honavar SG, Singh AD. Experience with eyelid-sparing orbital exenteration: the 2000 Tullos O. Coston Lecture. *Ophthalmic plastic and reconstructive surgery*. 2001;17(5):355-61.
8. Looi A, Kazim M, Cortes M, Rootman J. Orbital reconstruction after eyelid- and conjunctiva-sparing orbital exenteration. *Ophthalmic plastic and reconstructive surgery*. 2006;22(1):1-6.
9. Yeatts RP. The esthetics of orbital exenteration. *American journal of ophthalmology*. 2005;139(1):152-3.
10. Rodrigues ML, Kohler HF, Faria JC, Ikeda MK, Vartanian JG, Kowalski LP. Reconstruction after extended orbital exenteration using a fronto-lateral flap. *International journal of oral and maxillofacial surgery*. 2009;38(8):850-4.
11. Suarez C, Ferlito A, Lund VJ, Silver CE, Fagan JJ, Rodrigo JP, et al. Management of the orbit in malignant sinonasal tumors. *Head & neck*. 2008;30(2):242-50.
12. Croce A, Moretti A, D'Agostino L, Zingariello P. Orbital exenteration in elderly patients: personal experience. *Acta otorhinolaryngologica Italica : organo ufficiale della Societa italiana di otorinolaringologia e chirurgia cervico-facciale*. 2008;28(4):193-9.
13. Ackuaku-Dogbe E. Review of orbital exenterations in Korle-Bu teaching hospital. *Ghana medical journal*. 2011;45(2):45-9.
14. Frunza A, Slavescu D, Zamfirescu D, Stanculescu L, Grintescu I, Enache V, et al. Orbital exenteration - a salvage procedure? *Romanian journal of morphology and embryology = Revue roumaine de morphologie et embryologie*. 2013;54(4):1161-7.
15. Hoffman GR, Jefferson ND, Reid CB, Eisenberg RL. Orbital Exenteration to Manage Infiltrative Sinonasal, Orbital Adnexal, and Cutaneous Malignancies Provides Acceptable Survival Outcomes: An Institutional Review, Literature Review, and Meta-Analysis. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2015.
16. Kaur A, Khattri M, Jaiswal V. Pattern of orbital exenteration at a tertiary eye care centre in North India. *Nepalese journal of ophthalmology : a biannual peer-reviewed academic journal of the Nepal Ophthalmic Society : NEPJOPH*. 2012;4(1):64-7.
17. Maheshwari R. Review of orbital exenteration from an eye care centre in Western India. *Orbit*. 2010;29(1):35-8.
18. Nassab RS, Thomas SS, Murray D. Orbital exenteration for advanced periorbital skin cancers: 20 years experience. *Journal of plastic, reconstructive & aesthetic surgery : JPRAS*. 2007;60(10):1103-9.
19. Qasemyar A, Aljudaibi N, Wavreille O, Mortier L, Martinot-Duquennoy V, Guerreschi P. Orbital exenteration and periorbital skin cancers. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons*. 2014;72(4):811-6.
20. Soysal HG. Orbital exenteration: a 10-year experience of a general oncology hospital. *Orbit*. 2010;29(3):135-9.
21. Ben Simon GJ, Schwarcz RM, Douglas R, Fiaschetti D, McCann JD, Goldberg RA. Orbital exenteration: one size does not fit all. *American journal of ophthalmology*. 2005;139(1):11-7.

22. Rahman I, Cook AE, Leatherbarrow B. Orbital exenteration: a 13 year Manchester experience. *The British journal of ophthalmology*. 2005;89(10):1335-40.
23. Ali MJ, Pujari A, Dave TV, Kaliki S, Naik MN. Clinicopathological profile of orbital exenteration: 14 years of experience from a tertiary eye care center in South India. *International ophthalmology*. 2015.
24. Atique-Tacla M, Paves L, Pereira MD, Manso PG. [Exenteration: a retrospective study]. *Arquivos brasileiros de oftalmologia*. 2006;69(5):679-82. Exenteracao: estudo retrospectivo.
25. Levin PS, Dutton JJ. A 20-year series of orbital exenteration. *American journal of ophthalmology*. 1991;112(5):496-501.
26. Mohr C, Esser J. Orbital exenteration: surgical and reconstructive strategies. *Graefe's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 1997;235(5):288-95.
27. Naquin HA. Exenteration of the orbit. *AMA archives of ophthalmology*. 1954;51(6):850-62.
28. Rathbun JE, Beard C, Quickert MH. Evaluation of 48 cases of orbital exenteration. *American journal of ophthalmology*. 1971;72(1):191-9.
29. Simons JN, Robinson DW, Masters FW. Malignant tumors of the orbit and periorbital structures treated by exenteration. *Plastic and reconstructive surgery*. 1966;37(2):100-4.
30. Putterman AM. Orbital exenteration with spontaneous granulation. *Archives of ophthalmology*. 1986;104(1):139-40.
31. Levin PS, Ellis DS, Stewart WB, Toth BA. Orbital exenteration. The reconstructive ladder. *Ophthalmic plastic and reconstructive surgery*. 1991;7(2):84-92.
32. Reese AB. Exenteration of the orbit; with transplantation of the temporalis muscle. *American journal of ophthalmology*. 1958;45(3):386-90.
33. Reese AB, Jones IS. Exenteration of the orbit and repair by transplantation of the temporalis muscle. *American journal of ophthalmology*. 1961;51:217-27.
34. Dortzbach RK, Hawes MJ. Midline forehead flap in reconstructive procedures of the eyelids and exenterated socket. *Ophthalmic surgery*. 1981;12(4):257-68.
35. Shore JW, Burks R, Leone CR, Jr., McCord CD, Jr. Dermis-fat graft for orbital reconstruction after subtotal exenteration. *American journal of ophthalmology*. 1986;102(2):228-36.
36. Mauriello JA, Jr., Han KH, Wolfe R. Use of autogenous split-thickness dermal graft for reconstruction of the lining of the exenterated orbit. *American journal of ophthalmology*. 1985;100(3):465-7.
37. Shields JA, Shields CL, Suvarnamani C, Tantisira M, Shah P. Orbital exenteration with eyelid sparing: indications, technique, and results. *Ophthalmic surgery*. 1991;22(5):292-7.

# **EXPANSIVE HYDROGEL PELLETS IN THE TREATMENT OF POST ENUCLEATION/EVSCERATION SOCKET SYNDROME**

Delmas Juliette, Lossouran Adrien, Adenis Jean-Paul, Robert Pierre-Yves

1. Centre Hospitalier Universitaire Dupuytren – Limoges - France

## **Goal**

To evaluate the post-operative result after orbital injection of expansive hydrogel pellets, in the treatment of enophthalmos secondary to PESS.

## **Materials and methods**

This study was carried out between January 2010 and September 2015, in the University Hospital Dupuytren Limoges – France. Twenty surgeries were performed in 16 patients (20 procedures). Intra-orbital injection of the hydrogel pellets was performed under general anesthesia, behind the hydroxyapatite orbital implant, through small conjunctival incisions (1.5 mm). The volume of a single hydrogel pellet volume is 0.024 ml, and it increases tenfold after intra-orbital rehydration, after 24 to 48 hours.

## **Results**

The average enophthalmos rose from 13.35 +/- 3.69 (6-19) mm pre-injection to 14.98 +/- 3.69 (7-20) post-injection. The average number of injected pellets was 7.45 +/- 2.76 (3-15) per procedure. Three sockets (18.75%) needed a second surgery, with the same procedure. Esthetic result after fitting a new prosthetic device was satisfactory in 100% of the cases. Half of the patients felt a post operative pain, successfully treated by analgesic. One of our patient presented a sub-cutaneous implant migration, although not requiring surgical correction.

## **Conclusions**

This procedure appears sure, safe, quick, without risk of any serious complication, and leading to good results. It can be performed on ambulatory surgery. The expected post-operative increase of enophthalmos is 1 to 3 mm. Up to 10 or 11 pellets can be injected but more may lead to severe post-operative pain. Long term results seem to stable over time but a longer follow-up is necessary to evaluate the risk of late implant migration or extrusion.

## **Bibliography**

- Tyers AG, Collin JR. Orbital implants and post enucleation socket syndrome. *Trans Ophthalmol Soc U K.* 1982 Apr;102 (Pt 1):90–2.
- Wiese KG, Vogel M, Guthoff R, Gundlach KK. Treatment of congenital anophthalmos with self-inflating polymer expanders: a new method. *J Cranio-Maxillo-fac Surg Off Publ Eur Assoc Cranio-Maxillo-fac Surg.* 1999 Apr;27(2):72–6.
- Gundlach KKH, Guthoff RF, Hingst VHM, Schittkowski MP, Bier UC. Expansion of the socket and orbit for congenital clinical anophthalmia. *Plast Reconstr Surg.* 2005 Oct;116(5):1214–22.



## **OUR EXPERIENCE WITH AN AUTOMATED PHOTO-OPHTHALMIC DEVICE FOR MEASURING MRD**

Simon Barnard<sup>1</sup>, Giuseppe Reppucci<sup>2</sup>

1. Barnard Levit Optometrists, London, UK
2. Ascon Contactlinsen, Deutschland

### **Goal**

To provide comments and observations on a new automated method of measuring MRD.

### **Materials and methods**

The photo-ophthalmic device described in this presentation is used routinely in the author's practice to measure external eye data. This data is included when referring patients to oculoplastic surgeons for ptosis or lid retraction evaluation.

### **Results**

An example of a patient with lid retraction is presented to describe the data obtained with the device.

### **Conclusions**

Our experience is that the device measures MRD 1 & MRD 2 in patients with partial ptosis or lid retraction providing objective documentation. An independent jpeg photo with a grid scale provides an additional measuring scheme when ptosis obscures the Purkinje image.

### **Bibliography**

- Rosen C, Ramdass D, Norman C, Buckingham R. (2015) Visible Iris Diameter with the Volk Eye Check Device, Poster, BCLA conference UK
- Ayoub T, Hamed S, Uddin J (2016) Our experience of the Volk Eye Check for lid measurements, International Thyroid Eye Disease Symposium London, UK

## **SURGICAL MANAGEMENT OF RESIDUAL BROW SUSPENSION PTOSIS AND SEVERE LAGOPHTALMOS**

Dr.Michel Tazartes, Dr.Sorinela Roata, Dr. Saad Mouine

- National Center Hospital of Ophthalmology CHNO des XVXX,Paris, France

### **Goal**

Surgical approach of residual ptosis with severe lagophtalmos and corneal scar previously operated 10 years before with frontalis sling is described. We emphasize careful preop case examination of levator function, sling suspension action with superior eyelid entropion, corneal exposure and scarring and looking for uncorrected congenital ectropion often associated with congenital ptosis.

### **Materials and methods**

Bilateral procedure was performed under general anesthesia. We repositioned the autogenous fascia lata sling and adjusted the superior eyelid position in order to obtain the best symmetrical eyelid contour. Tarsal incisions helped evert the superior eyelid. To correct lagophtalmos, we performed retractors resection and reinsertion by conjunctival approach. Same approach was used for a malar lift and canthopexy in a conjugated effort to correct the lagophtalmus and the aspect of “ droopy face “ of this young patient

### **Results**

Both anatomically and functionally results were satisfactory with patient very happy with the esthetic results.Lagophtalmos was corrected and the eyelid contour was judged as Appropriate.

### **Conclusions**

Congenital ptosis surgical correction can sometimes be very tricky because of general anesthesia and the unpredictable results of frontalis sling technique. Careful preop examination looking for any congenital ectropion has to be done in order to correct both ptosis and ectropion otherwise eyelid elevation could complicate with lagophtalmos. If possible, associating a malar lift would help restore the eyelid closure and function.

### **Bibliography**

- J.Nerad, Techniques in Ophtalmic Plastic Surgery
- A.Tyers, R. Collin, Ophtalmic Plastic Surgery
- C.McCord, Frontalis suspension, Eyelid surgery principles and techniques
- C.McCord, Evaluation of the ptosis patient, Eyelid surgery principles and techniques

## **ORBITAL AND ADNEXA INVOLVEMENT IN ANGIOLYMPHOID HYPERPLASIA WITH EOSINOPHILIA**

Alicia Galindo-Ferreiro, Hind M Alkatan, Sahar Elkhamary, Azza Maktabi, Silvana Schellini, Antonio Augusto Cruz, Patrica Akaishi

1. King Khaled Eye Specialist Hospital, Saudi Arabia
2. Department of Ophthalmology, Complejo Asistencial de Palencia, Palencia, Spain
3. Department of Ophthalmology, Otorhinolaryngology and Head and Neck Surgery, School of Medicine of Ribeirão Preto–University of São Paulo, Brazil
4. Department of Ophthalmology Faculdade de Medicina de Botucatu - UNESP, São Paulo, Brasil

### **Goal**

Angiolymphoid Hyperplasia with Eosinophilia (ALHE) is an unusual benign disorder and a very rare condition in the orbit and ocular adnexa. We are presenting demographic data from 7 cases of ALHE with computed tomography (CT), magnetic resonance (MR) images and typical histopathological confirmation.

### **Case series**

The patients were attended between 2000 to 2015, at King Khaled Eye Specialist Hospital, Saudi Arabia and School of Medicine of Ribeirão Preto, Brazil. None of these patients developed systemic lymphoma during the follow-up. All except one case had good resolution with treatment.

### **Bibliography**

- Acocella, A., C. Catelani and P. Nardi (2005). "Angiolymphoid hyperplasia with eosinophilia: a case report of orbital involvement." *J Oral Maxillofac Surg* 63(1): 140-144.
- Alder, B., A. Proia and J. Liss (2013). "Distinct, bilateral epithelioid hemangioma of the orbit." *Orbit* 32(1): 51-53.
- Archer, K. F., J. J. Hurwitz and G. Heathcote (1991). "Orbital angiolymphoid hyperplasia with eosinophilia. Presentation as chalazion." *Ophthal Plast Reconstr Surg* 7(3): 208-221.
- Azari, A. A., M. R. Kanavi, M. Lucarelli, V. Lee, A. M. Lundin, H. D. Potter and D. M. Albert (2014). "Angiolymphoid hyperplasia with eosinophilia of the orbit and ocular adnexa: report of 5 cases." *JAMA Ophthalmol* 132(5): 633-636.
- Baker, M. S., R. B. Avery, C. R. Johnson and R. C. Allen (2012). "Methotrexate as an alternative treatment for orbital angiolymphoid hyperplasia with eosinophilia." *Orbit* 31(5): 324-326.
- Bangal, S., R. Chitgopekar, A. Gupta and R. Karle (2011). "Orbital extension of supraorbital angiolymphoid hyperplasia with eosinophilia." *Australas Med J* 4(3): 111-113.
- Barakova, D., J. Sach, P. Kuchynka, M. Redinova and I. Kocur (2002). "[Angiolymphoid hyperplasia with eosinophilia with bilateral involvement of the lacrimal glands]." *Klin Monbl Augenheilkd* 219(5): 376-379.
- Bostad, L. and W. Pettersen (1982). "Angiolymphoid hyperplasia with eosinophilia involving the orbita. A case report." *Acta Ophthalmol (Copenh)* 60(3): 419-426.
- Buggage, R. R., C. W. Spraul, T. H. Wojno and H. E. Grossniklaus (1999). "Kimura disease of the orbit and ocular adnexa." *Surv Ophthalmol* 44(1): 79-91.

- Chun, S. I. and H. G. Ji (1992). "Kimura's disease and angiolymphoid hyperplasia with eosinophilia: clinical and histopathologic differences." J Am Acad Dermatol 27(6 Pt 1): 954-958.
- Cook, H. T. and N. D. Stafford (1988). "Angiolymphoid hyperplasia with eosinophilia involving the lacrimal gland: case report." Br J Ophthalmol 72(9): 710-712.
- Cunniffe, G., T. Alonso, C. Dinares, F. J. Medina and R. Medel (2014). "Angiolymphoid hyperplasia with eosinophilia of the eyelid and orbit: the Western cousin of Kimura's disease?" Int Ophthalmol 34(1): 107-110.
- Esmaili, D. D., E. L. Chang, T. M. O'Hearn, R. E. Smith and N. A. Rao (2008). "Simultaneous presentation of Kimura disease and angiolymphoid hyperplasia with eosinophilia." Ophthalm Plast Reconstr Surg 24(4): 310-311.
- Fernandes, B. F., A. Al-Mujaini, T. Petrogiannis-Halioitis, A. Al-Kandari, B. Arthurs and M. N. Burnier, Jr. (2007). "Epithelioid hemangioma (angiolymphoid hyperplasia with eosinophilia) of the orbit: a case report." J Med Case Rep 1: 30.
- Gonzalez-Cuyar, L. F., F. Tavora, X. F. Zhao, G. Wang, A. Auerbach, N. Aguilera and A. P. Burke (2008). "Angiolymphoid hyperplasia with eosinophilia developing in a patient with history of peripheral T-cell lymphoma: evidence for multicentric T-cell lymphoproliferative process." Diagn Pathol 3: 22.
- Hidayat, A. A., J. D. Cameron, R. L. Font and L. E. Zimmerman (1983). "Angiolymphoid hyperplasia with eosinophilia (Kimura's disease) of the orbit and ocular adnexa." Am J Ophthalmol 96(2): 176-189.
- Jang, K. A., S. J. Ahn, J. H. Choi, K. J. Sung, K. C. Moon, J. K. Koh and Y. H. Shim (2001). "Polymerase chain reaction (PCR) for human herpesvirus 8 and heteroduplex PCR for clonality assessment in angiolymphoid hyperplasia with eosinophilia and Kimura's disease." J Cutan Pathol 28(7): 363-367.
- Kempf, W., A. C. Haeffner, K. Zepter, C. A. Sander, M. J. Flaig, B. Mueller, R. G. Panizzon, T. Hardmeier, V. Adams and G. Burg (2002). "Angiolymphoid hyperplasia with eosinophilia: evidence for a T-cell lymphoproliferative origin." Hum Pathol 33(10): 1023-1029.
- Krema, H. and N. El-Bolkainy (2014). "A stepwise multimodality treatment of diffuse angiolymphoid hyperplasia of the orbit." Orbit 33(1): 75-77.
- Lin, B., S. H. Tan and A. Looi (2008). "Angiolymphoid hyperplasia with eosinophilia of the eyelid with spontaneous regression." Ophthalm Plast Reconstr Surg 24(4): 308-310.
- McEachren, T. M., S. Brownstein, D. R. Jordan, V. A. Montpetit and R. L. Font (2000). "Epithelioid hemangioma of the orbit." Ophthalmology 107(4): 806-810.
- Moss, H. B., D. T. Sines, J. Blatt, J. J. Dutton and A. D. Proia (2012). "Epithelioid hemangioma responsive to oral propranolol." Ophthalm Plast Reconstr Surg 28(4): e88-90.
- Sanchez-Acosta, A., D. Moreno-Arredondo, R. I. Rubio-Solornio, H. A. Rodriguez-Martinez and A. A. Rodriguez-Reyes (2008). "Angiolymphoid hyperplasia with eosinophilia of the lacrimal gland: a case report." Orbit 27(3): 195-198.
- Sanchez-Orgaz, M., A. Insausti-Garcia, L. Y. Gregorio, A. A. Duralde and R. Romero-Martin (2014). "Epithelioid hemangioma of the orbit or angiolymphoid hyperplasia with eosinophilia." Ophthalm Plast Reconstr Surg 30(3): e70-72.
- Sheren, S. B., P. L. Custer and M. E. Smith (1989). "Angiolymphoid hyperplasia with eosinophilia of the orbit associated with obstructive airway disease." Am J Ophthalmol 108(2): 167-169.
- Shields, C., J. Shields and R. Glass (1990). "Bilateral orbital involvement with angiolymphoid hyperplasia with eosinophilia (Kimura's disease)." Press 9(2): 89-95.

- Wolff, D., H. Andree, I. Hilgendorf, J. Casper, M. Freund and C. Junghanss (2008). "Sirolimus in combination with tacrolimus in allogeneic stem cell transplantation--timing and conditioning regimen may be crucial." Biol Blood Marrow Transplant 14(8): 942-943.

## **ORBITAL TUMOR IN CHILDHOOD, UN UNSPECTED PRESENTATION**

Avila Jorge Ezequiel

- Hospital de Clinicas “Jose de San Martin” Buenos Aires, Argentina

### **Goal**

To analyze an unusual form of presentation from vascular malformation as an orbital tumor.

### **Materials and methods**

We present a case of a child of 1 year and 5 months old who consult for proptosis of 4 months ago. Personal history denied. It is noted an unpain proptosis, with EOM conserved, with no alterations on slit lamp and with normal fundus exam. Orbital CT reveals a diffuse infiltration of the intra and extraconal fat, without precise limits. We decide to program an incisional biopsy.

### **Results**

We start the biopsy entering to the orbit through the palpebral sulcus and take samples of the injury. Pathology informs a tumor with numerous vessels of thin walls and flats endothelium. Immunohistochemical reveals GLUT-1 negative, CD-34 positive and ALFA ACTINE OF SMOOTH MUSCLE positive, which confirm venous malformation. ISSVA classification 2014 (Vascular anomalies). They are slow-growing benign lesions, characterized by thin-walled sinusoids containing blood or hematic collections without apparent venous flow. They present from birth but there are visible weeks or months later. the incidence is 1.5%, 2/3 are venous. This malformations have low flow.

### **Conclusions**

Vascular malformations have varied presentations. The study of pathology as well immunohistochemical are fundamental to classify these diseases for proper treatment and monitoring them. At this moment, several treatments are known, like corticosteroids, propranolol, interferon, vincristine or timolo, being the orbital presentation very responders to therapy with corticosteroids.

### **Bibliography**

- Vascular malformations: classification and terminology the radiologist needs to know. Lowe LH<sup>1</sup>, Marchant TC, Rivard DC, Scherbel AJ. Semin Roentgenol. 2012 Apr;47(2):106-17. doi: 10.1053/j.ro.2011.11.002.
- Current management of hemangiomas and vascular malformations. Clinics in Plastic Surgery, Volume 32, Issue 1, Pages 99-116 Jennifer J. Marler, John B. Mulliken

- GLUT-1: an extra diagnostic tool to differentiate between haemangiomas and vascular malformations J. Leon-Villapalosa, K. Wolfeb, L. Kangesua, Department of Plastic and Reconstructive Surgery, St Andrews Centre for Plastic Surgery and Burns, Court Road, Broomfield, Chelmsford CM1 7ET, UK Department of Histopathology, Southend General Hospital, Prittlewell Chase, Westcliff-on-Sea, Essex SSO ORY, UK. British Journal Of Plastic Surgeons.

# SPONTANEOUS PREOPERATIVE RUPTURE OF DERMOID CYSTS: CLINICOPATHOLOGICAL AND RADIOLOGICAL FEATURES

Alicia Galindo-Ferreiro, Patricia Akaishi, Yerena Muiños-Diaz, Silvana Schellini, Augusto Cruz, Sahar Elkhamary, Hind Alkhatan, Rajiv Khandekar

5. Department of Oculoplastics, King Khaled Eye Specialist Hospital, Saudi Arabia
6. Department of Ophthalmology, Complejo Asistencial de Palencia, Palencia, Spain
7. Department Of Diagnostic Radiology, Mansoura Faculty Of Medicine, Egypt
8. Department of Ophthalmology, Otorhinolaryngology and Head and Neck Surgery, School of Medicine of Ribeirão Preto–University of São Paulo, Brazil
9. Research department, King Khaled Eye Specialist Hospital, Saudi Arabia
10. Department of Ophthalmology Faculdade de Medicina de Botucatu - UNESP, São Paulo, Brazil
11. Department of Ophthalmology, King Saud University College of Medicine, Riyadh, Saudi Arabia

## Goal

To evaluate the association of clinical signs of inflammation, computed tomography (CT) findings and histological features to determine if pre-rupture of orbital dermoid cysts can influence outcomes

## Materials and methods

A retrospective chart review was performed of patients with orbital dermoid cyst managed between 1989 to 2014. Data were collected on clinical features, CT scan and histology postoperatively to confirm signs of pre-rupture of the cyst. The associations were tested with Odd's ratio and 95% confidence interval (CI). Receiver operating characteristic (ROC) and the area under the curve was calculated to determine an association between cyst size and pre-rupture.  $P < 0.05$  was statistically significant.

## Results

Forty-two patients with dermoids were evaluated. The median age was 5 years. There were 23 (54.8%) patients with cysts located temporally and 19 (45.2%) with medial cysts. Thirty-six (85.7%) patients had a superficial cyst. Pre-rupture occurred in 61.9% of patients, with median age of 5.5 years old. Median age of patients in whom pre-rupture did not occur was 3 years old. Pre-rupture occurred more frequently in larger (median=16 mm) and medially located cysts. Bone scalloping was more frequent in pre-rupture cases (65.4%). Fat was the most common content for all cases, but, there was a greater association between liquid and pre-rupture ( $P=0.08$ ). The sensitivity of CT in predicting pre-surgical rupture was 50% and specificity was 75%. The size of the dermoid allowed accurate determination of pre-rupture in 72.5% of the cases. Histology confirmed pre-rupture in 61.9% of the cysts. Outcomes were not affected by intraoperative cyst rupture.



## Conclusions

There was a positive association between bony remodeling and histological pre-rupture. The age of the patient and size of the dermoid were related to signs of pre-rupture. Early surgical removal of dermoids is required to avoid pre-rupture. There was no relationship between pre-rupture and intraoperative complications

## Bibliography:

1. Abou-Rayyah Y, Rose GE, Konrad H, Chawla SJ, Moseley IF. Clinical, radiological and pathological examination of periocular dermoid cysts: evidence of inflammation from an early age. *Eye*. 2002;16(5):507-12.
2. Shields JA, Augsburger JJ, Donoso LA. Orbital dermoid cyst of conjunctival origin. *American journal of ophthalmology*. 1986;101(6):726-9.
3. Shields JA, Kaden IH, Eagle RC, Jr., Shields CL. Orbital dermoid cysts: clinicopathologic correlations, classification, and management. The 1997 Josephine E. Schueler Lecture. *Ophthalmic plastic and reconstructive surgery*. 1997;13(4):265-76.
4. Colombo F, Holbach LM, Naumann GO. Chronic inflammation in dermoid cysts: a clinicopathologic study of 115 patients. *Orbit*. 2000;19(2):97-107.
5. Bonavolonta G, Tranfa F, de Conciliis C, Strianese D. Dermoid cysts: 16-year survey. *Ophthalmic plastic and reconstructive surgery*. 1995;11(3):187-92.
6. Eibl KH, Kampik A, Hintschich C. Upper eyelid oedema from a dumbbell-shaped dermoid cyst. *Acta ophthalmologica Scandinavica*. 2005;83(1):126-7.
7. Fasina O, Ogun OG. Giant deep orbital dermoid cyst presenting early in infancy in a Nigerian child: a case report and review of the literature. *Journal of medical case reports*. 2012;6:320.
8. Nugent RA, Lapointe JS, Rootman J, Robertson WD, Graeb DA. Orbital dermoids: features on CT. *Radiology*. 1987;165(2):475-8.
9. Sherman RP, Rootman J, Lapointe JS. Orbital dermoids: clinical presentation and management. *The British journal of ophthalmology*. 1984;68(9):642-52.
10. Bonavolonta G, Strianese D, Grassi P, Comune C, Tranfa F, Uccello G, et al. An analysis of 2,480 space-occupying lesions of the orbit from 1976 to 2011. *Ophthalmic plastic and reconstructive surgery*. 2013;29(2):79-86.
11. Dizon M, Ozturk A, Redett RJ, Izbudak I. Neuroimaging findings in a child with dumbbell-shaped frontosphenoidal dermoid cyst presenting as preseptal cellulitis. *Pediatric radiology*. 2009;39(8):850-3.
12. Burgener F, Korman M, Pudas T. *Differential Diagnosis in Conventional Radiology*. 2008.
13. Ray MJ, Barnett DW, Snipes GJ, Layton KF, Opatowsky MJ. Ruptured intracranial dermoid cyst. *Proceedings*. 2012;25(1):23-5.
14. Berg JP, Barrett RT, Young AK, Marx DP. Orbital dermoid in a patient with bilateral anophthalmia: a case and review of the literature. *Orbit*. 2014;33(6):475-6.
15. del Rocio Arce Gonzalez M, Navas A, Haber A, Ramirez-Luquin T, Graue-Hernandez EO. Ocular dermoids: 116 consecutive cases. *Eye & contact lens*. 2013;39(2):188-91.
16. Das CJ, Tahir M, Debnath J, Pangtey GS. Neurological picture. Ruptured intracranial dermoid. *Journal of neurology, neurosurgery, and psychiatry*. 2007;78(6):624-5.
17. Shields JA, Shields CL. Orbital cysts of childhood--classification, clinical features, and management. *Survey of ophthalmology*. 2004;49(3):281-99.

18. Cavazza S, Laffi GL, Lodi L, Gasparrini E, Tassinari G. Orbital dermoid cyst of childhood: clinical pathologic findings, classification and management. *International ophthalmology*. 2011;31(2):93-7.
19. Ahuja R, Azar NF. Orbital dermoids in children. *Seminars in ophthalmology*. 2006;21(3):207-11.
20. Akbaba M, Karsloglu S, Damlack A, Karcoglu ZA. Intradiploic epidermoid cyst of frontal bone with spontaneous fistulization to eyelid. *Ophthalmic plastic and reconstructive surgery*. 2012;28(1):e15-7.
21. Lane CM, Ehrlich WW, Wright JE. Orbital dermoid cyst. *Eye*. 1987;1 ( Pt 4):504-11.
22. Emerick GT, Shields CL, Shields JA, Eagle RC, Jr., De Potter P, Markowitz GI. Chewing-induced visual impairment from a dumbbell dermoid cyst. *Ophthalmic plastic and reconstructive surgery*. 1997;13(1):57-61.
23. Missotten GS, van Santbrink-Bakker H, de Keizer RJ. Dumbbell-shaped dermoid cysts. *Acta ophthalmologica*. 2010;88(2):e11-2.
24. Min LC. CT and UltraSound in the diagnosis of orbital dermoid cysts – A retrospective study at the University Teaching Hospital in Lusaka. *Medical Journal of Zambia*.35(2):58-61.
25. Sathananthan N, Moseley IF, Rose GE, Wright JE. The frequency and clinical significance of bone involvement in outer canthus dermoid cysts. *The British journal of ophthalmology*. 1993;77(12):789-94.

## **ENDOSCOPIC PRIMARY DACRYOCYSTORHINOSTOMY: ARE SILICONE TUBES NEEDED?**

M. Cavaliere, M.D.; G. Concilio\*, M.D.; F.M. Folino\*, M.D.; M. Iemma, M.D.  
Department Otorhinolaryngology, University Hospital “Ruggi”, Salerno, Italy.  
\* Department of Ophthalmology, University Hospital “Ruggi”, Salerno, Italy.

### **Goal**

The first dacryocystorhinostomy (DCR) was described by Caldwell in 1893 but only 100 years later with the development of modern nasal endoscopes the endoscopic DCR has become popular and it's considered the gold standard treatment for patients with obstruction of the lacrimal system on the level of the sac or below it.

The most common reasons for failure in endoscopic DCR are adhesions and restenosis. Bicanalicular insetion of a silicone tube into the lacrimal duct is the most common procedure to prevent the closure of the rhinostomy. However, the silicone tube itself may cause tissue granulation predisposing the site to infection and adhesions. The objective of this study is to assess different outcomes between endoscopic dacryocystorhinostomy (En-DCR) with and without silicone intubation.

### **Methods and materials**

We retrospectively analyzed 40 patients (46 procedures), suffering from chronic epiphora for primary acquired nasolacrimal duct obstruction, treated between January 2013 and June 2015 with En-DCR and divided into two groups depending on silicone stent intubation.

The surgical outcomes were evaluated using Munk's score criteria. Functional success was defined as absence of epiphora, no further episodes of dacryocystitis, and a patent ostium after saline irrigation.

The surgery was performed under general anesthesia in all cases.

Step 1. Mucosal incision (Fig. 1 ) - The mucosal incisions were made with scalpel scythed. The incision was started approximately 5-6 mm above the insertion of the middle turbinate and carried anteriorly for approximately 8mm.

Step 2. Flap elevation - A suction elevator was used to raise the mucosa from the underlying bone along the maxillary line posteriorly to the thin lacrimal bone.

Step 3. Bone removal - The thin lacrimal bone was dissected free of the posterior aspect of the lacrimal sac. The thick bone of the frontal process of the maxilla was initially removed using a Kerrison rongeur and then continue with an angled diamond burr.

Step 4. Removal of the lacrimal sac mucosa - After the medial wall of the lacrimal sac has been exposed, sac was tented by the lacrimal probe (Fig. 2). The medial wall was incised vertically (Fig. 3) and angled Blakesley forceps were used to remove the redundant tissue along the incised sac edges.

Step 5. Intubation (Fig. 4) – In 20 procedures lacrimal probes threaded with Silastic tubes are passed through the canaliculi, directed out of the nasal cavity, and tied.

## **Results**

20 En-DCR with stent and 26 En-DCR without stent were performed. The period of the insertion of silicone tube was from 3 months to 6 months, and average  $4.1 \pm 1.2$  months. The follow-up observation period was from 11 months to 48 months, and average  $26.2 \pm 5.4$  months.

Success rate were, respectively, 90% in the stent group and 88.5 % in the non-stent group with no statistical differences (Student's test). The ostial size reduction has been reported in higher percentage in the stent group, mainly due to peristomal granuloma and scar tissue formation.

## **Discussion**

En-DCR without silicone stent intubation should be the first choice of procedure, stent intubation should be reserved in selected cases with poor local conditions pre and intra-operatively assessed (tight space, too intraoperative trauma). The benefits of non-intubation are less patient discomfort, reduced surgical time and costs, simpler follow-up regimen and less intubation-associated complications (granuloma, scar tissue, etc.).

## **Bibliography**

- Chen S, Le CH, Liang J. Practice patterns in endoscopic dacryocystorhinostomy: survey of the American Rhinologic Society. *Int Forum Allergy Rhinol*. 2016 [Epub ahead of print]
- Heichel J, Hassan T, Bredehorn-Mayr T, Wienke A, Struck HG. [External Dacryocystorhinostomy--Analysis of Patient Material of the University Hospital Halle from 2000 to 2011]. *Klin Monbl Augenheilkd*. 2016;233(1):29-37.
- Jung SK, Kim YC, Cho WK, Paik JS, Yang SW. Surgical outcomes of endoscopic dacryocystorhinostomy: analysis of 1083 consecutive cases. *Can J Ophthalmol*. 2015;50(6):466-70.

## **RESULTS OF PROBING UNDER ENDOSCOPIC GUIDANCE IN CHILDREN FORTY-EIGHT MONTHS AND OLDER**

Silvana Schellini MD, Alicia Galindo-Ferreiro, Rajiv Khandekar, Patricia Mitiko Akaishi, Augusto Cruz

1. King Khaled Eye Specialist Hospital, Saudi Arabia
2. Department of Ophthalmology, Complejo Asistencial de Palencia, Palencia, Spain
3. Department of Ophthalmology, Otorhinolaryngology and Head and Neck Surgery, School of Medicine of Ribeirão Preto–University of São Paulo, Brazil
4. Department of Ophthalmology Faculdade de Medicina de Botucatu - UNESP, São Paulo, Brasil

### **Goal**

This study evaluates the success rates of endoscopy assisted probing or conventional probing in children  $\geq 48$  months old.

### **Materials and methods**

This retrospective study included children 48 months or older with CNLDO who underwent endoscopic assisted probing or conventional probing between January 2011 to August 2015 at a tertiary eye care hospital in central Saudi Arabia. Probing was considered successful when symptoms disappeared (subjective success) and dye disappearance test was normal (objective success). Demographic data, clinical features, intraoperative and postoperative variables were correlated to the success rate.

### **Results**

One hundred and twelve children with CNLDO undergoing endoscopic assisted (37 patients) or conventional (75 patients) probing were included. The objective and subjective success rates of endoscopic assisted and conventional probing was 94.6% (95% CI 89.5- 99.7) and 58.7% (95% CI 47.6 – 69.8) respectively. The success rate was higher in the endoscopic group and the difference favored the endoscopic procedure in older children.

### **Conclusions**

Endoscopic assisted probing had significantly higher success rates than conventional probing in children 48 months or older. Endoscopic assisted probing can achieve good outcomes to treat CNLDO even in children over 48 months old.

### **Bibliography:**

1. Choi WC, Kim KS, Park TK, Chung CS: Intranasal endoscopic diagnosis and treatment in congenital nasolacrimal duct obstruction. *Ophthalmic surgery and lasers*, 33(4) (2002) 288-292.
2. Kouri AS, Tsakanikos M, Linardos E, Nikolaidou G, Psarommatis I: Results of endoscopic assisted probing for congenital nasolacrimal duct obstruction in older children. *International journal of pediatric otorhinolaryngology.*, 72(6) (2008) 891-896.
3. MacEwen CJ, Young JD, Barras CW, Ram B, White PS: Value of nasal endoscopy and probing in the diagnosis and management of children with congenital epiphora. *The British journal of ophthalmology*. 85(3) (2001) 314-318.

4. Wallace EJ, Cox A, White P, Macewen CJ: Endoscopic-assisted probing for congenital nasolacrimal duct obstruction. *Eye*. 20(9) (2006) 998-1003.
5. Al-Faky YH: Nasal endoscopy in the management of congenital nasolacrimal duct obstruction. *Saudi J Ophthalmol*.28(1) (2004)6-11.
6. Robb RM: Success rates of nasolacrimal duct probing at time intervals after 1 year of age. *Ophthalmology*.105(7) (1998) 1307-1309; discussion 1309-1310.
7. Zwaan J: Treatment of congenital nasolacrimal duct obstruction before and after the age of 1 year. *Ophthalmic surgery and lasers*. 28(11) (1997)932-936.
8. el-Mansoury J, Calhoun JH, Nelson LB, Harley RD: Results of late probing for congenital nasolacrimal duct obstruction. *Ophthalmology*. 93(8) (1986) 1052-1054.
9. Ciftci F, Akman A, Sonmez M, Unal M, Gungor A, Yaylali V: Systematic, combined treatment approach to nasolacrimal duct obstruction in different age groups. *European journal of ophthalmology*.10(4) (2000) 324-329.
10. Alañón-Fernández MÁ, Alañón-Fernández FJ, Martínez-Fernández A, del Mar Górgora M, Calero B, López-Marín I, Alarcón S. Comparative study of primary intention lacrimal probing with and without nasal endoscopy. *Acta Otorrinolaringologica (English Edition)*. 2014;65(5):297-301.
11. Dean AG SK, Soe MM: *OpenEpi: Open Source Epidemiologic Statistics for Public Health*. 2015.
12. Al-Faky YH, Mousa A, Kalantan H, Al-Otaibi A, Alodan H, Alsuhaibani AH: A prospective, randomised comparison of probing versus bicanalicular silastic intubation for congenital nasolacrimal duct obstruction. *The British journal of ophthalmology*. 99(2) (2015) 246-250.
13. Elmorsy S, Shabana YK, Fayek HM: Endoscopic assisted probing for symptomatic congenital nasolacrimal duct obstruction after one year of age. *Rhinology* 2010, 48(1) (2010) 100-103.
14. Maheshwari R: Success rate and cause of failure for late probing for congenital nasolacrimal duct obstruction. *Journal of pediatric ophthalmology and strabismus*.45(3) (2008)168-171.
15. MacEwen CJ YJ: Epiphora in the first year of life. *Eye*. 1991, 5 (1991) 596-600.
16. Arora S, Koushan K, Harvey JT: Success rates of primary probing for congenital nasolacrimal duct obstruction in children. *Journal of AAPOS : the official publication of the American Association for Pediatric Ophthalmology and Strabismus / American Association for Pediatric Ophthalmology and Strabismus*.16(2) (2012) 173-176.
17. Casady DR, Meyer DR, Simon JW, Stasior GO, Zobal-Ratner JL: Stepwise treatment paradigm for congenital nasolacrimal duct obstruction. *Ophthalmic plastic and reconstructive surgery*. 22(4) (2006) 243-247.
18. Frick KD, Hariharan L, Repka MX, Chandler D, Melia BM, Beck RW, Pediatric Eye Disease Investigator G: Cost-effectiveness of 2 approaches to managing nasolacrimal duct obstruction in infants: the importance of the spontaneous resolution rate. *Archives of ophthalmology*.129(5) (2011) 603-609.
19. Eshragi B, Fard MA, Masomian B, Akbari M: Probing for congenital nasolacrimal duct obstruction in older children. *Middle East African journal of ophthalmolog*. 20(4) (2013) 349-352.
20. Hung CH, Chen YC, Lin SL, Chen WL: Nasolacrimal Duct Probing under Topical Anesthesia for Congenital Nasolacrimal Duct Obstruction in Taiwan. *Pediatrics and neonatology* 2015.
21. Isaza G, Arora S: Probing without irrigation in children with congenital nasolacrimal duct obstruction. *Clinical and investigative medicine Medecine clinique et experimentale*. 36(3) (2013) E158-162.
22. Miller AM, Chandler DL, Repka MX, Hoover DL, Lee KA, Melia M, Rychwalski PJ, Silbert DI, Pediatric Eye Disease Investigator G, Beck RW et al: Office probing for treatment of

- nasolacrimal duct obstruction in infants. *Journal of AAPOS : the official publication of the American Association for Pediatric Ophthalmology and Strabismus / American Association for Pediatric Ophthalmology and Strabismus*.18(1) (2014) 26-30.
23. Pediatric Eye Disease Investigator G, Repka MX, Chandler DL, Beck RW, Crouch ER, 3rd, Donahue S, Holmes JM, Lee K, Melia M, Quinn GE et al: Primary treatment of nasolacrimal duct obstruction with probing in children younger than 4 years. *Ophthalmology*.115(3) (2008) 577-584 e573.
  24. Zilelioglu G, Hosal BM: The results of late probing in congenital nasolacrimal duct obstruction. *Orbit*.26(1) (2007) 1-3.
  25. Sasaki H, Takano T, Murakami A: Direct endoscopic probing for congenital lacrimal duct obstruction. *Clinical & experimental ophthalmology*.41(8) (2013)729-734.
  26. Theodoropoulou S, Sutherland MS, Haddow K, Blaikie A: Success rates of endoscopic-assisted probing for congenital nasolacrimal duct obstruction in children. *The Journal of laryngology and otology*.127(8) (2013) 794-798.
  27. Yagci A, Karci B, Ergezen F: Probing and bicanalicular silicone tube intubation under nasal endoscopy in congenital nasolacrimal duct obstruction. *Ophthalmic plastic and reconstructive surgery* 2000, 16(1) (2000) 58-61.
  28. Cakmak SS, Yildirim M, Sakalar YB, Keklikeci U, Alakus F: Is it necessary to accompany probing with endoscopy in cases of congenital nasolacrimal canal obstruction? *International journal of pediatric otorhinolaryngology*.74(9) (2010) 1013-1015.
  29. Al-Faky YH, Al-Sobaie N, Mousa A, Al-Odan H, Al-Huthail R, Osman E, Al-Mosallam AR: Evaluation of treatment modalities and prognostic factors in children with congenital nasolacrimal duct obstruction. *Journal of AAPOS : the official publication of the American Association for Pediatric Ophthalmology and Strabismus / American Association for Pediatric Ophthalmology and Strabismus*.16(1) (2012) 53-57.
  30. Mannor GE, Rose GE, Frimpong-Ansah K, Ezra E: Factors affecting the success of nasolacrimal duct probing for congenital nasolacrimal duct obstruction. *American journal of ophthalmology*.127(5) (1999) 616-617.
  31. Kashkouli MB, Beigi B, Parvareh MM, Kassaei A, Tabatabaee Z: Late and very late initial probing for congenital nasolacrimal duct obstruction: what is the cause of failure? *The British journal of ophthalmology*. 87(9) (2003) 1151-1153.
  32. Leone CR, Jr., Van Gemert JV: The success rate of silicone intubation in congenital lacrimal obstruction. *Ophthalmic surgery*.21(2) (1990)90-92.
  33. Schellini SA, Narikawa S, Ribeiro SC, Nakagima V, Padovani CR, Padovani CR: [Congenital lacrimal obstruction: outcome and factors associated with therapeutic probing]. *Arquivos brasileiros de oftalmologia*.68(5) (2005) 627-630.

# TEMPORARY MEDIAL UPPER EYELID LAGOPHTHALMOS AFTER EXTERNAL DACRYOCYSTORHINOSTOMY

Ivan O. Haefliger, M.D., FEBO<sup>1,2</sup>, Otmar Meienberg, M.D.<sup>2,3</sup>, and Ana Rosa Pimentel de Figueiredo, M.D., Ph.D.<sup>4</sup>

<sup>1</sup> Holbein-Praxis *KLINIK*, Holbeinstrasse 29, 4051 Basel Switzerland,

<sup>2</sup> School of Medicine, University of Basel, Klingenbergstrasse 61, 4056 Basel, Switzerland,

<sup>3</sup> Neurologie am Schaulager, Emil Frey-Strasse 85, 4142 Münchenstein, Switzerland

<sup>4</sup> Department of Oculoplastic Surgery, São Geraldo Hospital, Federal University of Minas Gerais, Av. Prof. Alfredo Balena, 190, Santa Efigênia Belo Horizonte - MG, 30130-100, Brazil (Chairman: Ana Rosa Pimentel de Figueiredo)

## Goal

Report of three cases of medial upper eyelid lagophthalmos as complication of external dacryocystorhinostomy.

## Materials and methods

Shortly after dacryocystorhinostomy (side of the nose skin incision), three patients ( $28 \pm 4$  years; mean  $\pm$  standard deviation), out of ten consecutive ones, presented with an ipsilateral lagophthalmos of  $4 \pm 1$  mm by voluntary eyelid closure and  $6 \pm 1$  mm by spontaneous blink. The lagophthalmos appeared to primarily result of a medial upper eyelid paresis. Patient 1 complained bitterly of dry eye symptoms and of her lagophthalmos. Patient 2 had mild symptoms but got very concerned after peers made her aware of an asymmetric blink. Patient 3 was asymptomatic and did not notice anything particular.

## Results

Lagophthalmos resolved spontaneously within three months after surgery, first by recovery of voluntary eyelid closure and then of spontaneous blinking.

## Conclusions

Temporary lagophthalmos can occur as a complication of external dacryocystorhinostomy, most likely due to damage of the only recently described superficial buccal and/or zygomatic branches of the facial nerve that in subjects run upward to cross over the medial ligament and innervate the medial part of the orbicularis oculii muscle.

## Bibliography

- Haefliger IO, Meienberg O, Pimentel de Figueiredo AR. Temporary Medial Upper Eyelid Lagophthalmos after External Dacryocystorhinostomy. *Klin Monbl Augenheilkd*. 2016 Apr;233(4):406-8.



## **DIFFERENTS FORMS OF CONJUNCTIVAL AND EYELID LYMPHOMAS**

Camacho Belén, Ignacio Racana, Tula Romina, Billagra Alejandra, Weil Daniel

1. Hospital de Clínicas Jose de San Martin, Buenos Aires, Argentina

### **Goal**

Presentation of cases of lymphoma in eyelid and conjunctiva in patients and different presentation start

### **Materials and methods**

Review of clinical history multidisciplinary photographic and imaging documentation and review of literature

### **Results**

Patient of 53 years who consulted for palpebral tumor in the right lower eyelid of 6 months of evolution, with personal history of major bilateral BTE adenomegalia the contralateral side. It presents the ophthalmological examination: visual acuity of 8/10 in both eyes without correction, 10/10 correction, bmc: od temporary scleritis, conjunctival fornix mass and two lower eyelid lesions; oi unremarkable, MOE preserved without diplopia. Fundus: no particularities. Computed Tomography in hyperdense lesion was evident in scleral insertion of the lateral rectus. excisional biopsy and tarsal conjunctival lesion and cancer screening was performed. flow cytometry was performed, confirming the diagnosis with biopsy result conjunctival and orbital NHL lymphoma type MALT lineage. It was indicated marrow aspiration and chemotherapy for 6 months indicated. Patient of 65-year-old who consults for red eye of 6 months duration, no history, treated as viral conjunctivitis, visual acuity of 10/10 in both eyes without correccion, bmc: conjunctival fornix mass on od, rest without particularities. Computed tomography in no pathologies are observed; excisional biopsy confirmed diagnosis of lymphoma Conjunctival type of NHL, MALT lineage is performed. Patient 60 years old, who consults for unilateral proptosis of 3 months of evolution, that the examination had av sc 10/10 without correction, bmc: unremarkable, HERTEL 22 mm od, oi 17mm, fo: no particularities. Computed Tomography in hyperdense lesion was evident in superointerna retro-orbital area. It is performed biopsy confirmed diagnosis of MALT-cell NHL

### **Conclusions**

Lymphomatous lesions conjunctival lesions are presented as salmon-colored sacks fund but in some cases may occur conmitantes associated with orbital lesions with infiltration of muscles extraoculares. En ophthalmologic examination must analyze the conjunctiva and funds sac and contralateral eyelid. with excisional biopsy diagnosis is made and treatment is excision plus radiation therapy and or quemioterapia.

**Bibliography:**

- Auw-Haedrich C1, Coupland SE, Kapp A, Schmitt-Gräff A, Buchen R, Witschel H. Long term outcome of ocular adnexal lymphoma subtyped according to the REAL classification. Revised European and American Lymphoma; University Eye Hospital, Freiburg, Germany. Br J Ophthalmol. 2001 Jan;85(1):63-9.
  
- Rath S1, Connors JM, Dolman PJ, Rootman J, Rootman DB, White VA Comparison of American Joint Committee on Cancer TNM-based staging system (7th edition) and Ann Arbor classification for predicting outcome in ocular adnexal lymphoma.;Department of Ophthalmology and Visual Sciences , University of British Columbia, Vancouver, Canada. Orbit. 2014 Feb;33(1):23-8.. Epub 2013 Nov 1.
- White WL, Ferry JA, Harris NL, Grove AS., Jr Ocular adnexal lymphoma. A clinicopathologic study with identification of lymphomas of mucosa-associated lymphoid tissue type. Ophthalmology. 1995 Dec;102(12):1994–2006.

## **SEBACEOUS CARCINOMA OF THE EYELID WITH PAGETOID INFILTRATION OF THE CONJUNCTIVA: TWO CASE REPORTS**

Ricardo Díaz Céspedes, Gloria Gornals Montés, Diamar Pardo Lopez, Enrique España Gregori.David Salom Alonso.

- Hospital de Manises, Valencia; España

### **Goal**

To show the importance of Sebaceous Gland Carcinoma (SGC) of the eyelid. It is a very rare, slow growing, malignant tumor. Most ocular sebaceous carcinomas originate in the tarsal meibomian glands. It commonly occurs in the sixth to eighth decade of life, with preponderance for women and Asians. The upper eyelid is affected more frequently. The disease can spread directly by orbital invasion, lymphatic spread to regional lymph nodes and hematogenous spread to distant organs. Correct diagnosis of SGC is often delayed due to its resemblance to a variety of benign conditions of the eyelids such as chalazion or chronic blepharconjunctivitis.

### **Materials and methods**

A review of the literature was performed, focusing on the epidemiology, clinical presentation, differential diagnosis, treatment and monitoring

### **Results**

We report on two female patients suffering from sebaceous carcinoma of the upper lid with pagetoid infiltration of the conjunctiva. The time span from initial treatment up to definite diagnosis and treatment was one year . In both cases the tumor was radically removed surgically. Histopathologically, it were diagnosed as a SGC. There was no lymphovascular invasion and no metastasis; therefore no other treatment was included.

### **Conclusions**

SGC is a rare but aggressive neoplasm that tends to clinically and histopathologically mimic other conditions. Remains a common diagnostic pitfall for both the clinician and histopathologist. Therapy of choice is the radical surgical removal of the tumor. It should be considered in cases of persistence of conjunctivitis or chalazion despite appropriate therapeutic interventions.

### **Bibliography**

- Prieto-Granada C, Rodriguez-Waitkus P. Sebaceous Carcinoma of the Eyelid. *Cancer Control*. 2016 Apr;23(2):126-32.
- Pe'er J. Pathology of eyelid tumors. *Indian J Ophthalmol*. 2016 Mar;64(3):177-90. doi: 10.4103/0301-4738.181752.
- Aravindan KP. Sebaceous carcinoma. *Indian J Pathol Microbiol*. 2015 Jul-Sep;58(3):408, 415. doi: 10.4103/0377-4929.162937.

- Kaliki S, Ayyar A, Dave TV, Ali MJ, Mishra DK, Naik MN. Sebaceous gland carcinoma of the eyelid: clinicopathological features and outcome in Asian Indians. *Eye (Lond)*. 2015 Jul;29(7):958-63. doi: 10.1038/eye.2015.79. Epub 2015 May 22.
- Mulay K, White VA, Shah SJ, Honavar SG. Sebaceous carcinoma: clinicopathologic features and diagnostic role of immunohistochemistry (including androgen receptor). *Can J Ophthalmol*. 2014 Aug;49(4):326-32. doi: 10.1016/j.jcjo.2014.04.004. Epub 2014 Jul 19.
- Mulay K, Aggarwal E, White VA. Periocular sebaceous gland carcinoma: A comprehensive review. *Saudi J Ophthalmol*. 2013 Jul;27(3):159-65. doi: 10.1016/j.sjopt.2013.05.002.
- Rene C. Oculoplastic aspects of ocular oncology. *Eye (Lond)*. 2013 Feb;27(2):199-207. doi: 10.1038/eye.2012.243. Epub 2012 Nov 30.
- Esmali, Bitar et al. American Joint Committee on Cancer T Category for Eyelid Sebaceous Carcinoma Correlates with Nodal Metastasis and Survival. *Ophthalmology* 2012;119:1078-1082.
- Deprez, Manuel and Uffer, Sylvie. Clinicopathological Features of Eyelid Skin Tumors. A Retrospective Study of 5504 Cases and Review of Literature. *Am J Dermatopathol* 2009; 31:256-262.
- Buitrago, William and Aaron Joseph. Sebaceous carcinoma: the great masquerader. *Dermatologic Therapy*. Vol 21, 2008, 459-466.
- Dores GM, Curtis RE, Toro JR, Devesa SS, Fraumeni JF Jr. Incidence of cutaneous sebaceous carcinoma and risk of associated neoplasms: insight into Muir-Torre syndrome. *Cancer*. 2008;113(12):3372-81
- Song A, Carter KD, Syed NA, Song J, Nerad JA. Sebaceous cell carcinoma of the ocular adnexa: clinical presentations, histopathology, and outcomes. *Ophthal Plast Reconstr Surg*. 2008;24(3):194-200.
- Shields JA, Demirci H, Marr BP, Eagle RC Jr, Shields CL. Sebaceous carcinoma of the eyelids: personal experience with 60 cases. *Ophthalmology*. 2004;111(12):2151-7.

# **TOTAL RECONSTRUCTION FUNCTIONAL OF THE UPPER AND LOWER EYELID USING COMBINED FLAPS AND FREE GRAFTS**

Torres Ana Marina. MD (1) Dolmetsch Angela M, MD (2)  
(1) Instituto de Diagnóstico de La Plata, La Plata, Argentina  
(2) Clínica de Oftalmología de Cali, Cali, Colombia

## **Goal**

To describe a surgical technique for total reconstruction of the upper eyelid and the lower eyelid in two cases using combined flaps and free skin and mucosal grafts.

## **Materials and methods**

Retrospective and descriptive study of two patients with advanced malignant tumor recurrence (one of them with Sebaceous cell carcinoma and one with Basal Cell carcinoma). The patients had total reconstruction of the upper eyelid and partial reconstruction of the lower eyelid with the use of three different types of flaps (Periostium, Tripiet and Fricke) and free skin and mucosal grafts.

## **Results**

Surgery produced complete reconstruction of upper and lower eyelids in both cases with good functional and cosmetic results. There was no tumor recurrence and the follow up time was 18 months.

## **Conclusions**

Total reconstruction of the upper eyelid and partial reconstruction of the lower eyelid can be performed satisfactorily with good functional and cosmetic results with the use of combined flaps and free skin and mucosal grafts. In this way adequate eyelid function and vision can be preserved.

## **Bibliography**

- Jean-Louis deSousa, Franzco; Igal Leibovitch, MD; Raman Malhotra, FRCOphth; Brett O'Donnell, Franzco; Tim Sullivan, Franzco; Dinesh Selva, Fracs, Franzco. Techniques and Outcomes of Total Upper and Lower Eyelid Reconstruction. Arch Ophthalmol. 2007;125(12):1601-1609. doi:10.1001/archophth.125.12.1601.
- Bertrand, Baptiste M.D.; Colson, Thomas Roger Jr.; Baptista, Claire M.D.; Georgiou, Charalambos M.D.; Philandrianos, Cécile M.D., Ph.D.; Degardin, Nathalie M.D., Ph.D.; Bardot, Jacques M.D.; Casanova, Dominique M.D. Total Upper and Lower Eyelid Reconstruction: A Rare Procedure—A Report of Two Cases. Plastic & Reconstructive Surgery: October 2015;136(4): 855-859

# **TRANSCONJUNCTIVAL FAT REPOSITION WITHOUT PEDICLE IN LOWER BLEPHAROPLASTY**

Torres, A. Marina

- Instituto de Diagnóstico de La Plata

## **Goal**

To present and describe the experience of the author in lower eyelid blepharoplasty, a surgical technique transconjunctival fat transfer without pedicle, in the nasojugal fold (tear trough) in order to achieve an aesthetic improvement.

## **Materials and methods**

Retrospective, observational and descriptive study where clinical history was reviewed and images for a period of 2 years. They were excluded from the study patients with a history of eyelid surgery and / or lifting of the middle third of the face. Described in detail surgical technique, representative photographs are presented to assess the outcome; the results and complications were reviewed.

## **Results**

They were included in the study 14 patients. The average age was 62 years (range 49-75) and mean follow-up was 6 months (range 2-13). It was included 12 women and 2 men. No postoperative complications were recorded. All patients were satisfied with the surgical results felt.

## **Conclusions**

The technique of fat repositioning without pedicle provides a subtle change in the contour of the lower eyelid. It is not as powerful as alloplastic implant, which other author have suggested, but has the advantage to be autogenous tissue; or to have used the lipofilling. In the author's experience has shown subjectively a decrease in depth fold nasojugal, is used filler patient's own fat and the same area. No longer requiring additional time or specific fat processing. The complications are the same for any other variation of transconjunctival blepharoplasty.

## **Bibliography**

- Goldberg RA, Edelstein C, Shorr N (1999) Fat repositioning in lower blepharoplasty to maintain infraorbital rim contour. *Facial Plast Surg* 15:225–229
- Nassif PS (2005) Lower blepharoplasty: transconjunctival fat repositioning. *Facial Plast Surg Clin North Am* 13:553–559
- Loeb R (1993) Nasojugal groove leveling with fat tissue. *Clin Plast Surg* 20:393–400
- Coleman SR (1997) Facial contouring with liposculpture. *Clin Plast Surg* 24:347–367

## **ORBITAL MUCOCELE**

Arce.Jessica, Avila.Jorge, Billagra.Alejandra, Camacho. Belén, Gasparini.Cecilia, Magnano. Paula, Racana. Ignacio, Tula. Romina, Weil. Daniel

- Hospital de Clínicas José de San Martín. CABA. Buenos Aires. Argentina

### **Goals**

The aim of our poster is to show four patients with different types of mucocele of paranasal sinus and their signs and symptoms.

### **Materials and methods**

We present four cases of orbital mucocele, their locations and their signs and symptoms.

### **Results**

Dermoid and epidermoid cystic, angiofibromas, neurofibromas, osseous fibromas, cylindromas, inverted papillomas, cholesterol granulomas, and odontogenic cysts may cause expansion similar to mucoceles in the sinus walls and, therefore, they should be included in the differential diagnosis. CT is the most important and beneficial method, and it reflects the imaging of homogeneous, cystic, expansile mass eroding the surrounding bone tissue. MRI is used to determine the involvement of neighboring soft tissue and to distinguish the lesion from other soft tissue neoplasms. Mucoceles are treated surgically. The location, magnitude, and expansion of the lesion are determinants of the appropriate surgical procedure. The endoscopic approach is a faster procedure with less morbidity and a shorter time. One exception is when the frontal sinus is compromised laterally, in that case the best approach is an orbitotomy.

### **Conclusions**

Mucocele is a rare condition that can affect any paranasal sinus. Therefore it is not a well-known disease for the general ophthalmologist. The location, clinical manifestations, magnitude and expansion are determinants of the surgical approach.

### **Bibliography**

- D.Weil: Mucocele orbitario. Diagnóstico y tratamiento. Experiencia en 19 casos. *An. Inst Barraquer*. 26: 17-30 (1997)
- JJ Khong, R Malhotra, PJ Wormald and D Selva: Endoscopic sinus surgery for paranasal sinus mucocele with orbital involvement. *Eye* (2004) 18, 877-881.
- Har-El G. Endoscopic management of 108 sinus mucoceles. *Laryngoscope* 2001, 111:2131-2134.
- Murat Topdag, Mete Iseri, Fatih Sari, Selvet Erdogan, I Gurkan Keskin: Paranasal sinus mucoceles: our clinical experiments. *Int J Clin Exp Med* 2015;8(10):18518-18522. –
- Hssaine K, Belhoucha B, Rochdi Y, Nouri H, Aderdour L, Raji A. Paranasal sinus mucoceles: About 32 cases. *Rev Stomatol Chir Maxillofac Chir Orale*. 2016 Feb;117(1):11-4.

## **ORBITAL HEMANGIOPERICYTOMA**

Cynthia Anagua, Cecilia Gasparini, Sabrina Bagdadi, Martin Rojo, Daniel Weil.

2. Home institution: Hospital de Clínicas de Buenos Aires.

### **Goal**

The goal of our poster is to report a case of Orbital Hemangiopericytoma we received in the Hospital, to describe both clinical and histopathological findings and the treatment of choice.

### **Materials and methods**

Clinical and histopathological features were reviewed in a 50 year old female patient who was confirmed as having primary orbital hemangiopericytoma in Hospital de Clínicas de Buenos Aires.

### **Results**

The patient who was diagnosed as having primary orbital hemangiopericytoma by histopathologic and clinical findings, presented with progressive proptosis and some associated symptoms such as extraocular motility limitation. Visual Field showed superior temporal scotoma. Neuroimaging showed a well-circumscribed mass in the superior medial aspect of the right orbit, a lateral displacement of the optical nerve, which compromised medial rectus and extended to the ethmoidal sinus and CNS. Biopsy was taken and later, due to its retro-orbital localization partial surgical resection was performed. At this time, radiotherapy is being considered as an adjuvant treatment.

### **Conclusions**

Both clinical and histopathological findings are important for diagnosing and evaluating the treatment outcomes of orbital hemangiopericytoma. Total excision is suggested as the treatment of choice, and radiotherapy can be an effective adjuvant treatment for recurrence.

### **Bibliography**

- Hsu CH, Wei YH, Peng Y, Liao SL. Orbital hemangiopericytoma in an Asian population. *J Formos Med Assoc.* 2014 Jun;113(6):356-63.
- Croxatto O., Font R., MD. "Tumors of the eye and ocular adnexa" *Atlas of Tumor Pathology.* Washington, DC.2006;5: 280-281.
- Ide F, Obara K, Mishima K, Saito I, Kusama K. Ultrastructural spectrum of solitary fibrous tumor: a unique perivascular tumor with alternative lines of differentiation. *Virchows Arch.* 2005;446:646–652.
- Sullivan TJ, Wright JE, Wulc AE, Garner A, Moseley I, Sathanathan N. Haemangiopericytoma of the orbit. *Aust N Z J Ophthalmol.* 1992 Nov;20(4):325-32.
- Croxatto O., Font R. Hemangiopericytoma of the orbit: a clinicopathologic study of 30 cases. *Hum patol.*1982;13: 210-218.
- Setzkorn RK Lee D Hemangiopericytoma of the orbit treated with conservative surgery and radiotherapy. *Arch Ophthalmol* 1987;105:1103-5.



- Shields JA, Shields Carol. "Atlas of Orbital Tumors". Philadelphia: WB Saunders, 1989:132-134.
- Henderson JW, Farrow GM: Primary Orbital hemangiopericytoma: an aggressive and potentially malignant neoplasm. Arch Ophthalmol 1978;96:666-673.

## **THE GARDEN OF MINERVA**

BILLAGRA ALEJANDRA<sup>1</sup> WEIL DANIEL<sup>1</sup>, COLANGELO LUIGI<sup>2</sup>

1.HOSPITAL DE CLNICAS JOSE DE SN MARTIN, CABA. ARGENTINA

2.UNIVERSITA DE CHIETI-CASA DI CURA SALUS.

### **Goal**

In the XXI century it survives over the years a small garden, overlooking the Mediterranean, surrounded by trees and flowers, is a faithful representative of the Medical School of Salerno from the Middle Ages Considered the first botanical garden, past show and present of this important area of medical and pharmacological studies.

### **Research Methodology**

"Ostensio simplicium" in the Simple Jardin del Maestro Mateo silvático, cultivated some of the plants from which the active ingredients used for therapeutic purposes in medicine of the time were obtained. In the master's work the silvático Pandette (*Opus pandectarum Medicinae*), with a simple lexicon 500 plastas describes your benefits medicinales..El I silvático work: of the 487 chapters on plants, 233 (42.9%) are defined with a name of Arab origin, 134 (27.6%) with a Greek origin and only 120 episodes (24.6%) are called by a Latin word

### **Materials and methods**

Different photographic shots of the city of Salerno and the Garden of Minerva, currently performed. Information was collected in museum and in bibliographic.

### **Results**

School was largely based on the Humoral Theory of Hippocrates and Galen, but also on the experience gained in the daily exercise. In addition, translations of Arabic texts, this experience is added a vast knowledge of herbal medicine and pharmacology. The characteristics of the Garden of Minerva and knowledge of pharmacology and botany of its founder Matthew silvático, made this space would allow the development of different plants, shrubs and tubers, but so also the scientific knowledge of physicians of the period.

### **Conclusions**

The Jardin de la Minerva has a special microclimate favored by the low incidence of the north wind and the favorable exposure, which, even today, allows the cultivation of plant species demanding in terms of humidity and heat, When walking through internal trails Garden not only see a rescued archaeological site, like many, but it represents an important link in the history of medicine and one of the first universities.

### **Bibliography**

- Pérgola F., Okner O., Historia de la Medicina. 1986.

- Fondazione Scuola Medica Salernitana. Giardino della Minerva Orto Botanico della Scuola Medica Salernitana. 2013.
- Cantarella E. Vitali A., Trotula de'Rugiero L'Armonia delle donne, trattato medievale di cosmesi con consiglio praticisul trucco e la cura del corpo.Ed. Manni. 2014

## UPPER EYELID RETRACTION IN GRAVE'S DISEASE

- Magnano Paula, Billagra Alejandra Avila Jorge, Racana Ignacio, Tula Romina, Weil Daniel  
- Hospital de Clínicas José de San Martín. CABA. Buenos Aires. Argentina

### Goals

The aim of our poster is to describe demographics, differential diagnosis, clinical manifestations and treatment methods of Upper eyelid retraction in thyroid eye disease

### Materials and methods

We present some patients corrected by different surgical techniques and the postoperative results.

### Results

Grave's upper eyelid retraction is a very frequent sign of Graves' orbitopathy which can be mild, moderate or severe. Between its differential diagnoses, contralateral ptosis, congenital retraction and Marcus Gunn phenomenon can be found.

Several lengthening technics have been proposed to correct this disorder, as müllerectomy, levator aponeurosis recession or levator aponeurosis transposition, being the most common complication hiper or hypocorrection. We present some cases treated by different techniques with good results.

### Conclusions

Upper eyelid retraction is a very frequent presentation in patients with Graves orbitopathy and many different approaches have been described to solve it. However, predictability of final eyelid height and contour following surgical correction remains problematic. We present some cases with good postoperative results using relatively easy and safe procedures.

### Bibliography

- Bartley GB (1996): The differential diagnosis and classification of eyelid retraction. *Ophthalmology* 103: 168–176.
- Ben Simon GJ, Mansury AM, Schwarcz RM, Modjtahedi S, McCann JD & Goldberg RA (2005): Transconjunctival Muller muscle recession with levator disinsertion for correction of eyelid retraction associated with thyroid-related orbitopathy. *Am J Ophthalmol* 140: 94–99.
- Harvey JT & Anderson RL (1981): The aponeurotic approach to eyelid retraction. *Ophthalmology* 88: 513–524.
- Mourits M, Koornneef L. Lid lengthening by sclera interposition for eyelid retraction in Graves' ophthalmopathy. *Br J Ophthalmol* 1991; 75 (6): 344–347

## **OPHTHALMOMYIASIS AND SKIN CANCER**

Billagra Alejandra, Matach Gustavo, Ortiz Basso Tomas

3. Hospital de Clinicas Jose de San Martin CABA, Hospital Nacional de Clinicas Cordoba, Hospital Nacional de Clinicas Santa Rosa, La Pampa.

### **Goal**

To report three cases of ocular myiasis related to oncologic lesions, review of the literature and pathology.

### **Materials and methods**

Three cases with medical records and their corresponding histopathology are presented. All patients are from Argentina and live in rural provinces of Argentina. They have been diagnosed and properly treated.

### **Results**

In the three cases, patients had infestation from larvae of *Cochliomya hominivorax* in ulcerated lesions of basal cell carcinoma. Neither manual larvae removal, nor antibiotic or antiparasitic treatment weren't enough to solve the orbital infestation. Orbit exenteration was needed to achieve final cure in all cases.

### **Conclusions**

Due to the potentially devastating consequences ending in mutilating surgery, it is mandatory to recognise and identify patients with risk of having myiasis infestation.

### **Bibliography**

- Hope FW. On insects and their larvae occasionally found in the human body. *Trans R Soc Entomol (Lond)* 1840; 2: 256–271.2.
- Raposo AA, Schettini AP, Massone C. Concurrent primary and secondary myiasis on basal cell carcinoma. *An Bras Dermatol.* 2012 Mar-Apr;87(2):292-5.3.
- Echchaoui A, Benyachou M, Houssa A, et al. A. Management of eyelid carcinomas: Retrospective bicentric study of 64 cases and review of the literature. *J Fr Ophtalmol.* 2016 Feb;39(2):187-94. Epub 2016 Jan 27. French.
- WHO. Classifications. International Classification of Diseases (ICD). MIASIS CIE-10 B87.2015.
- Robbins K, Khachemoune A. Cutaneous myiasis: a review of the common types of myiasis. *Int J Dermatol.* 2010 Oct; 49(10):1092-8. Review.
- Kersten RC, Shoukrey NM, Tabbara KF. Orbital myiasis. *Ophthalmology.* 1986 Sep;93(9):1228-32.
- Olea MS, Centeno N, Aybar CA, Ortega ES, et al. First Report of myiasis caused by *Cochliomyia hominivorax* (Diptera: Calliphoridae) in a diabetic foot ulcer patient in Argentina. *Korean J Parasitol Korean J Parasitol.* 2014 Feb;52(1):89-92. Epub 2014 Feb 19.

## **TREATMENT OF DRY EYE SYNDROME IN YOUNG AGE GLAUCOMA**

Claudia Guarracino<sup>1</sup>, Alessandra Trotta<sup>1</sup>, Giuseppe Maisto<sup>2</sup>

1 Università Federico II Napoli,

2 Università Torvergata di Roma

### **Goal**

This study born from the necessity of reducing symptoms of dry eye syndrome in patients subject to anti-glaucoma treatment. Thanks to the introduction of zinc sulphate and ialuronic acid, it has been opened a new path, more appropriate and less invasive by the chemical point of view.

### **Materials and methods**

This study, realized in 2016, controlled and randomized has been carried on 20 patients (40 eyes) that during the pre-surgical period has done:

BUT – SCHIRMER TEST – FERNING TEST – NO CONCACT TONOMETRY.

The tests has been repeated after 3 months using the same diagnostic instruments.

The chosen patients have instilled Zixol, single dose colllyirium 3 times a week for 3 months.

### **Results**

The test results show the real improvement of ocular surface.

### **Conclusions**

They've been analyzed the possible association of symptoms with the best results of lachrymal test and it has been noted an outstanding augmentation of BUT, as consequence of increased biolife of fat and mucinous molecules that holding water molecules, considerably increase corneal hydration, reducing symptomatology. Together with the more hydrating effect of ialuronic acid, it's possible to see reduction of symptoms linked to the foreign object's sensation.

### **Bibliography**

- De Paiva CS, Plugfelder SC. Tear Clearance Implication for ocular surface health. *Exp Eye Res* 2004; 78:395-397.
- Schiffman RM, Dale Christianson M, Jacobsen G, Hirsch JD, Reis BL. Reliability and validity of the Ocular Surface Disease Index (OSDI). *Arch Ophthalmol* 2000; 118: 615-21.
- Cook C, Foster P. Epidemiology of glaucoma: what's new? *Can J Ophthalmol* 2012;47:223-6.
- Weinreb RN, Zangwill LM, Jain S, et al. Predicting the onset of glaucoma: the confocal scanning laser ophthalmoscopy ancillary study to the Ocular Hypertension Treatment Study. *Ophthalmology*. 2010;117(9):1674-1683.
- Sullivan BD, Crews LA, Messmer EM, et al. Correlations between commonly used objective signs and symptoms for the diagnosis of dry eye disease: clinical implications. *Acta Ophthalmol* 2012 Dec 28

## **RIBOFLAVIN-BASED STAINING OF THE EPITHELIAL OCULAR SURFACE: A NEW DIAGNOSTIC OPPORTUNITY**

Salvatore Troisi, Ciro Caruso, Elisabetta Chiariello Vecchio, Decio Capobianco, Paola Vele, Salvatore Adriano Cirillo.

- Ospedale San Giovanni di Dio e Ruggi D'Aragona, Salerno

### **Purpose**

Our purpose was to evaluate the effectiveness and safety of a new dye solution with fluorescent properties containing Riboflavin (Droptest) for the slit lamp examination of the corneal and conjunctival epithelial defects and to compare with the traditional staining by fluorescein and lissamine green in Sjogren Syndrome patients.

### **Materials and methods**

We studied 20 eyes of 10 patients suffering from Sjogren Syndrome with bilateral epithelial corneal-conjunctival defects. Each patient was subjected to staining with Fluorescein solution and Lissamine green matches (F+L) in right eye and with Droptest solution (D) in left eye; after washing with NaCl 0,9% solution, we performed staining with D solution in right eye and with F+L in other eye. The observation was made by slit lamp with cobalt blue light after fluorescein administration and in white light after lissamine-dye; the examen after Droptest administration was made with cobalt blue light and anteposition of filter wratten #12 (yellow).

The evaluation was performed by two independent examiners for both eyes after each step, using the Lemp scheme, and underwent to statistic elaboration.

### **Results**

The average corneal score for the right eyes, based on Lemp scheme, was 4,2 using F+L staining and 4,1 after D coloration, whereas for the left eyes was 4,7 with F+L and 4,6 with D evaluation.

The average conjunctival score was in right eyes 5.1 for F+L and 5.0 for D, while in left eyes 5,4 for F+L and 5,4 for D.

No statistically significant difference between the two groups was found.

In 9 of 10 patients, after asking their preferences, Droptest© coloration was subjectively more tolerated than F+L procedure. No side effects reported.

### **Conclusions**

Our study highlights that this new riboflavin-solution technique (Droptest), combined with the anteposition of a wratten 12 filter on slit lamp, is equally effective and equally safety for the detection of corneal and conjunctival suffering areas, compared with the classic combined use of sodium fluorescein, specific for the corneal epithelium, and lyssamine green, specific for the conjunctival injuries. The riboflavin solution was preferred by 90% of the patients, with better patient compliance and reduction of examination time, compared with the evaluation using the white light for lyssamine and

cobalt blue light for fluorescein stain.. In conclusion the formulation containing riboflavine (Droptest) appears to represent an appropriate substitute of fluorescein in these type of patients, it is an innovative way, but further studies are needed.

## **Bibliography**

- 1) The architecture of the corneal stroma. Bron AJ. Br J Ophthalmol. 2001 Apr;85(4):379-81
- 2) Superficial punctate fluorescein staining of the ocular surface. Ward KW. Optom Vis Sci. 2008 Jan;85(1):8-16.
- 3) Rose bengal staining and cytologic characteristics associated with lipid tear deficiency. Lee SH, Tseng SC. Am J Ophthalmol. 1997 Dec;124(6):736-50.
- 4) Correlation between corneal sensitivity, subjective dry eye symptoms and corneal staining in Sjögren's syndrome. Adatia FA, Michaeli-Cohen A, Naor J, Caffery B, Bookman A, Slomovic A. Can J Ophthalmol. 2004 Dec;39(7):767-71
- 5) Practical double vital staining for ocular surface evaluation. Toda I, Tsubota K. Cornea. 1993 Jul;12(4):366-7
- 6) Patient tolerance and ocular surface staining characteristics of lissamine green versus rose bengal. Manning FJ, Wehrly SR, Foulks GN. Ophthalmology. 1995 Dec;102(12):1953-7
- 7) Optimizing evaluation of Lissamine Green parameters for ocular surface staining. Hamrah P, Alipour F, Jiang S, Sohn JH, Foulks GN. Eye (Lond). 2011 Nov;25(11):1429-34
- 8) An evaluation of the efficacy of fluorescein, rose bengal, lissamine green, and a new dye mixture for ocular surface staining. Korb DR, Herman JP, Finnemore VM, Exford JM, Blackie CA. Eye Contact Lens. 2008 Jan;34(1):61-4
- 9) ESCRS – European Society of cataract and refractive surgeons XXXIV Congress Copenhagen 10/09/2016 - 14/09/2016 A new biological coloration for corneal and conjunctival evaluation of epithelial injuries. S. Troisi, C. Caruso, M. Troisi, L. Pacente, E. Chiariello Vecchio, P. Cirillo.
- 10) An alternative to fluorescein for measuring intraocular pressure by means of Goldmann tonometer  
Maurizio Rolando, Francesca Bruzzone Ocular Surface & Dry Eye Center, ISPRE Oftalmica Istituto di Medicina Oftalmica, Genova, Italian Review of ophthalmology; 2,2,2016



## THE SCHOLA MEDICA SALERNITANA AND OPHTHALMOLOGY

aGiovanni Leo<sup>1</sup>, Rossella Russo<sup>2</sup>, Giusy Sbaglio<sup>3</sup>, Pasquale Attimonelli<sup>4</sup>, Luca Barile<sup>5</sup>, Orazio Campagna<sup>6</sup>

1.University of Naples Federico II, Italy

2.University of Salerno, Italy

3.Hospital De Luca and Rossano of Vico Equense, Italy

4.Hospital of Andria Operating Unit , Italy

5.Hospital Unit Umberto I of Nocera Inferiore, Italy

6.Sapienza University of Rome, Italy

### Purpose

The Schola Medica Salernitana dates back to Middle Ages. It was in strong contrast to the beliefs of that time and it became a pioneer institution to which the following universities would conform. Studies concerning the various branches of medicine would result in essays that eventually became true cornerstones. Our investigation, by tracing back the history of medicine, will allow us to rediscover the first approaches to eye surgery and the treatments which were considered most appropriate for the treatment of eye diseases.

### Investigation methodology

Strolling about the old town is not difficult to dive in institutional structures such as palaces, gardens, sculptures still relevant; the tradition is not supported by historical documents, dates the foundation of the ninth century school, through the work of four teachers: Helinus the Jew, Pontus the Greek, Adela the Arabic and Salernus the Latin. To relive the history of the Schola Medica Salernitana we went to the Virtual Museum of Salerno.

The first historical evidence of the activity of the school date back to the Thent century and are contained in the *Historia inventionis ac traslationis et miracula S. Trophimena*, in *Chronicon Hùgone of Flavigny*, and in the *Reims Richeiro Historia*.

Early surgical elements are found in the Thirteenth century, when in Salerno are studied the basic standards of teaching surgery, encoded by Ruggero from Frugardo and Rolando from Parma. It speaks for the first time of a surgery about eye diseases in the *Treaty of Ruggero from Frugardo Surgery*, who proposed treatments used to solve the red eyes and ocular hypertension proceeding with an engraving of intervention of the frontal artery.

### Results

The ophthalmic surgical practice was deeply analyzed in the second half of the XIII century thanks to the studies of Benvenuto Grafeo, author of *De Arte Probatissima Oculorum* and *De oculorum adfectibus*, which describes various ocular diseases and different surgical interventions. Another exponent of Salerno ophthalmic doctrine was David Armenio author of the *Tractatus de Oculis Accanamusali*, his work is considered the foundation of Salerno ophthalmology. We can find in this manuscript a very rare iconographic testimony documenting the eye surgical instruments of Salerno. In the

manuscript it is preserved a rare documentary iconography detailing eye surgery tools. The tools, depicted so clearly and precisely, are accompanied by a clear indication of their use: scissors, needles, hooks, chisels and cauteries for the removal of cataracts, lacrimal fistula, corneal blisters and other similar interventions.

### **Conclusions**

The medicine-eye surgery was one of the glories of the Schola Medica Salernitana. Thanks to the works of Benvenuto Graphaeo and David Armenio began important studies in this field which may be considered a reference point for all subsequent ophthalmological medicine.

### **Bibliography**

- De Renzi, *Storia documentata della Scuola Medica Salernitana*, Napoli, 1857
- Trad. G. Lauriello, *Post Mundi Fabricam manuale di Chirurgia*-Ed.Gaia 2011
- M. Pasca, *La Scuola Medica Salernitana*, Napoli,1988
- A. Musi, M.Oldoni, A. Placanica, *Storia dell'Università di Salerno*, Salerno,2001

## THE AESTHETIC MEDICINE IN THE SCHOLA MEDICA SALERNITANA: TROTULA DE RUGGIERO

Giovanni Leo\*, Rossella Russo\*\*, Valeria Russo\*\*\*, Lorena Manzur\*, Maria Cristiana Abate\*, Catello Russo\*

\* University of Naples Federico II, Italy    \*\*University of Salerno, Italy    \*\*\*Second University of Naples, Italy

### Object

The Schola Medica Salernitana was the first Cultural Center not controlled by the Church. It was the first and largest medical institution in Europe during the Middle Ages, therefore considered by many as the forerunner of modern universities. There began the translations from Arabic into Latin of medical texts from the ancient Greek scientists, making them accessible to Western scholars too. The school was also open to women who attended both as students and as teachers. Trotula is the most important exponent of the doctors who was interested in cosmetics. We can find her knowledge in one of the most important treatises of aesthetic medicine.

### Investigative methodology

The use of cosmetics to improve the human body is lost in the mists of time. Sources attest the use since prehistoric times, perhaps to strike fear at rivals and preys, for religious and magical rituals. The use of cosmetics, however, has changed during the time, following the concept of beauty, adapting to the taste, to the specific needs and trends.

The search for beauty and its origins led to Salerno, considered as the melting pot of Mediterranean civilization, which gave shape to this knowledge through the Schola Medica Salernitana and where took part many active women. We discovered visiting the Virtual Museum that among mulieres Salernitanae there was Abella, Rebecca Garner, Mercuriade, and last but not least, Trotula De Ruggiero (1050-1097) with his treatises *De Mulierum Passionanibus ante and post partum*, defined *Trotula major* concerning the gynecology and obstetrics and *De ornatu mulierum*, said *Trotula minor*, a treatise on skin diseases and their care.

### Results

Our journey back in the history of aesthetic medicine, has brought out an ancient treatise, "*De ornatu Mulierum*" which teaches women how to preserve and enhance their beauty and how to treat skin diseases through a series of precepts, tips and natural remedies. In the statement the author gives some make-up lessons, suggests how to hide wrinkles, remove puffiness from the face and eyes, shave the body, lighten the skin, hide blemishes and freckles, wash teeth and eliminate bad breath, dye hair, do the waxing, gingivitis and how to cure chapped lips. It also provides guidelines for preparing and using ointments and medicinal herbs for the face and hair and dispenses advices to improve wellbeing through steam baths and massages. Trotula describes in his treatise a primordial "scrub" for face. Indeed it recommends using an exfoliating

cleanser made with bread crumbs to smooth facial skin. It also describes a kind of primitive peeling using the irritant effect of the onion.

The paper reports 96 plants derivatives, 20 animal preparations and derivatives, 17 minerals, and 6 mixed preparations, such as ingredients for 63 total recipes, able to obtain as many remedies for cosmetic purposes and / or medicinal products.

### **Conclusions**

Trotula De Ruggiero had innovative ideas in many aspects: she considered that prevention was the main aspect of medicine and propagated new methods, unusual for the era, emphasizing the importance the hygiene, balanced nutrition and physical activity play for health. Most of the plants mentioned in the Treatise and also the other plants used in the school of Salerno for the experimental preparations, at the beginning, grew spontaneously in the area, then they were cultured together with other plants introduced in the Minerva Gardens of Salerno.

### **Bibliography**

- P. Cavallo Boggi, *Trotula e le malattie delle donne*, Torino, 1979.
- Ferraris, Z.A. and Ferraris, V.A. *Women of Salerno: contribution to the origins of surgery from medieval Italy*. Ann. Thorac. Surg. 1997.
- Green, M.H. *The Trotula. A Medieval Compendium of Women's Medicine*. Penn Press, Pennsylvania 2001.
- Cavallo P., Proto MC, Patruno C, Del Sorbo A, Bifulco M. *The first cosmetic treatise of history. A female point of view*, Int J Cosmet Sci. 2008





**Printed in May 2017**  
**Salerno - Italy**